



Pharmacological Effects of Garlic (*Allium sativum* L.)

Vinay K Singh, Dinesh K Singh*

Department of Zoology, DDU Gorakhpur University, Gorakhpur, U.P., INDIA

Received: 11 September 2007; accepted 21 January 2008

Online on 7 February 2008

Abstract

Singh VK, Singh DK. *Pharmacological Effects of Garlic (Allium sativum L.)*. *ARBS Annu Rev Biomed Sci* 2008;10:6-26. Garlic (*Allium sativum* L.) is a bulbous herb used as a food item, spice and medicine in different parts of the world. Its medicinal use is based on traditional experience passed from generation to generation. Researchers from various disciplines are now directing their efforts towards discovering the effects of garlic on human health. Interest in garlic among researchers, particularly those in medical profession, has stemmed from the search for a drug that has a broad-spectrum therapeutic effect with minimal toxicity. Recent studies indicate that garlic extract has antimicrobial activity against many genera of bacteria, fungi and viruses. The role of garlic in preventing cardiovascular disease has been acclaimed by several authors. Chemical constituents of garlic have been investigated for treatment of hyperlipidemia, hypertension, platelet aggregation and blood fibrinolytic activity. Experimental data indicate that garlic may have anticarcinogenic effect. Recent researches in the area of pest control show that garlic has strong insecticidal, nematocidal, rodenticidal and molluscicidal activity. Despite field trials and laboratory experiments on the pesticidal activity of garlic have been conducted, more studies on the way of delivery in environment and mode of action are still recommended for effective control of pest. Adverse effects of oral ingestion and topical exposure of garlic include body odor, allergic reactions, acceleration in the effects of anticoagulants and reduction in the efficacy of anti-AIDS drug Saquinavir.

©by São Paulo State University - ISSN 1806-8774

Keywords: garlic, allicin, alliin, cancer, hyperlipidemia, hypertension, antiplatelet activity, pesticide

Table of Contents

1. Introduction
2. Chemistry
3. Biological Effects
 - 3.1. Antimicrobial activities
 - 3.2. Garlic and cardiovascular diseases
 - 3.2.1. Hypocholesterolaemic action
 - 3.2.2. Hypotensive effect
 - 3.2.3. Antiplatelet aggregation activity
 - 3.3. Anticancer potential

* Correspondence

Dinesh K Singh. Reader, Department of Zoology, DDU Gorakhpur University, Gorakhpur-273 009, UP, India. E-mail: dksingh_gpu@yahoo.co.in

Vinay K Singh: vk Singh_gpu@yahoo.com

- 3.4. Garlic as a pesticide
 - 3.5. Adverse effects
 4. Conclusion
 5. Acknowledgments
 6. References
-

1. Introduction

Garlic, *Allium sativum* L. is a member of the Alliaceae family, has been widely recognized as a valuable spice and a popular remedy for various ailments and physiological disorders. The name garlic may have originated from the Celtic word 'all' meaning pungent. Cultivated practically throughout the world, garlic appears to have originated in central Asia and then spread to China, the Near East, and the Mediterranean region before moving west to Central and Southern Europe, Northern Africa (Egypt) and Mexico (Lutomski, 1987).

As one of the earliest cultivated plants, garlic is mentioned in the Bible and in the literature of Ancient Israel (The Talmud), Egypt (Codex Ebers) and India (Vedas and Purans, Charak Sanghita). Chinese strongly believe that garlic prolongs longevity (Srivastava *et al.*, 1995) and is useful in treating most human diseases, including infections, cancer and heart diseases. In historical writings of India and in literature like Vedas and Purans, garlic is reported to have medicinal as well as nutritive value in food items. Clinical indications of garlic include hemorrhoids, rheumatism, dermatitis, abdominal pain, cough, loss of appetite and loss of weight (Adetumbi & Lau, 1983; Anonymous, 1985; Koch & Lawson, 1996).

The medicinal qualities of garlic were described by Pliny the Elder, Aristophanes, and Galen. Hippocrates, the Father of Medicine, observed that garlic was excellent for curing tumors and is an effective diuretic. Aristotle attributed garlic as a cure for rabies, and the Prophet Mohammad recommended it for treating scorpion stings. In *Historia Naturalis* by Pliny (Fenwick & Hanley, 1985), garlic was recommended for gastro-intestinal disorders as well as dog and snake bites. Desired medicinal results of garlic are obtained when bulbs are chewed and swallowed or mixed with food and eaten.

Preparations of garlic are available as tablets, capsules, syrup, tinctures and oil. In ointment form, garlic has been used externally for treatment of ring worm; boiled with vinegar and sugar for treatment of asthma; made into an infusion for treatment of epilepsy; pounded with honey for use against rheumatism; and mixed with milk for use as a vermifuge. Garlic is commonly used in Europe and Asia for medicinal benefits in healing wounds, etc. (Bolton *et al.*, 1982; Srivastava *et al.*, 1995). In Germany, sale of garlic preparations competes with sales of leading drugs (Lawson, 1993).

Garlic produces a variety of volatile sulfur-based compounds which are effective as insect repellents and insecticides. Diallyl disulfide is one of such compounds which has a strong odor and acts as a powerful insecticide (Kaufman *et al.*, 1999). Commercial preparations of garlic are certified as insecticide against mites, nematodes and mosquito larvae affecting a variety of crops (Amonkar & Reeves, 1970; Gupta & Sharma, 1993; Gamboa-Leon *et al.*, 2006).

The objective of the present review is to highlight the pharmacological and therapeutic use of garlic in modern medicine and its use as pesticide against certain pest. Garlic chemistry follows in order to reach the aim of this review.

2. Chemistry

Intact garlic cloves contain only a few medicinally active compounds (Block, 1992; Lawson, 1993). The main chemical constituent of intact garlic is the amino acid alliin, an alkyl derivative of cysteine alkyl sulfoxide, which may vary from 0.2 to 2.0% fresh weight (Lutomski *et al.*, 1968; Kabelik, 1970; Michahelles, 1974; Lutomski, 1983a,b; 1987). The efficacy of chemical constituents of garlic chiefly depends on the mode of preparation of its extract. Crushing, chewing or cutting (or exposing dehydrated, pulverized garlic to water) of garlic cloves release the vacuolar enzyme allinase that rapidly lyses the cytosolic cysteine sulfoxide (Stoll & Seebeck, 1949a,b) to form sulfenic acid (R-SOH) (Block, 1992). Which immediately condenses to form allicin: the compound which produces the odor of fresh-cut garlic? Garlic contains at least 100 sulfur-containing compounds basic to medicinal uses (Lawson, 1993; Cai *et al.*, 1994; Srivastava *et al.*, 1995). Allicin represents 70-80 % of the total thiosulfonates

formed (Fig. 1). Other thiosulfinates formed include allyl-ss(o)-methyl (6-16% of total), methyl-ss(o)-allyl (3-9% of total), trans-1-propenyl (0.2-0.4% of total), trans-1-propenyl-ss(o)-methyl+methyl-ss(o)-trans-1-propenyl (0.1-2.5% of total) and methyl-ss(o)methyl (2% of total) thiosulfinates (Lawson *et al.*, 1991b). The formation of allicin (allyl 2-propene thiosulfinate or diallyl thiosulfinate) is complete in 0.2 to 0.5 min at room temperature. Formation of the methyl thiosulfinate takes 1.5 to 5.0 min (Lawson & Hughes, 1992). All possible combinations of 2-propene, 1-propene and methane sulfenic acids result in thiosulfinates (Lawson *et al.*, 1991a,b).




Figure 1. Conversion of alliin to allicin by the enzyme allinase, and of allicin into various sulphur-containing compounds.

The thiosulfinates released from crushed garlic are reactive molecules and undergo a number of transformations, depending on the temperature, pH and solvent conditions. Allyl-s- thiosulfinates (allicin) are the least stable of the eight thiosulfinates (Block, 1992; Lawson & Block, 1997). The half-life of allicin (concentration of 0.1-0.4mg/ml) at room temperature is 10 days in 1mM citric acid (pH 3), 4 days in water, 48 h in methanol or chloroform, 24 h in ethanol, 24 h in hexane and 3 h in ether (Lawson, 1993).

The principal transformation products after incubation of thiosulfinate in water are diallyl trisulfide, diallyl disulfide and allyl methyl trisulfide (Block, 1992). Incubation of allicin or allyl methane thiosulfinate in low polarity solvents (or without solvent) primarily produces 1, 3-vinyldithiin (2-vinyl-4H-, 3-dithiin) (51% of total), 1,2-vinyldithiin (3-vinyl-4H-1,2-dithiin) (19% of total) and lesser amounts of ajoene (E,Z-4,5,9-trithiododeca-1,6,11-trien 9-oxide) (12% of total) and sulfides (18% of total) (Fig. 1). Incubation of pure allicin in methanol yields 8% diallyl trisulfide and about equal amounts (20-25% of total) of ajoene and the vinyldithiins (Lawson *et al.*, 1991a; Lawson, 1993). Steam distillation of garlic bulbs yields a garlic oil that consists of diallyl (57% of oil), allyl methyl (37% of oil), and dimethyl (mono to hexa sulfides; 6% of oil of total) (Lawson, 1993). The essential oil content of fresh garlic bulbs is between 0.09 to 0.35 % (Lutomski, 1987).

3. Biological Effects

The biological activity of an extract of garlic depends on the mode of its preparation. The extract prepared at room temperature contains mainly allicin, which has a powerful antibacterial property and a strong smell. Besides allicin, small amounts of several other thiosulfinates and complex sulphanyl components, including the antithrombotic ajoenes, are also present. The enzyme allinase responsible for converting alliin to allicin is inactivated by heat. The water extract of heat-treated garlic mainly

contains alliin, which is odour-free. Garlic powder is a simply dehydrated, pulverized garlic clove. The allinase activity of garlic powder is identical to that of fresh garlic. It may be emphasized here that dehydration temperature should not exceed 60 °C, above which allinase is inactivated (Lawson, 1998). When distilled in steam, garlic yields an oily mass consisting of diallyl, methyl allyl, dimethyl, all originating from the thiosulfinates (Lawson *et al.*, 1991a; Lawson & Hughes, 1992). Garlic oil has been shown to possess biological properties such as antitumor and antioxidant effects (Sumiyoshi & Wargovich, 1989). Another widely studied garlic preparation is aged garlic extract (AGE). When garlic is subjected to cold aging process, organosulphur compounds, such as s-allyl cysteine, s-allyl mercaptocysteine, and several sulphur-containing amino acids, are produced. S-allyl cysteine and s-allyl mercaptocysteine have been shown to be anticarcinogenic and provides protection against liver damage (Srivastava *et al.*, 1995).

Garlic is one of the most investigated medicinal plants. During 1960 to 2007, more than three thousand research papers have been published on the chemistry and biological effects of garlic and garlic preparations. These studies mainly focus on the cardiovascular, anti-microbial and anti-cancer effects of garlic and, to a lesser extent, on the therapeutic indications for the treatment of hypoglycemia, heavy metal poisoning and liver dysfunction and hyperthyroidism (Lawson, 1993; Singh, 1995; Agarwal,

Table 1. Active constituents of garlic and their pharmacological actions.

Active component of garlic	Anti-biotic	Anti-fungal	Hyper-lipidemia	Anti-platelet aggregation	Anti-cancer	As a pesticide
alliin			+	+		
allicin	+	+	+	+	+	+
allixin					+	
adenosine				+	+	
Allyl1,5-hexadienyl trisulphide				+	+	
Allyl methyl trisulphide				+	+	
s-allyl2-pro pene thiosul phinate				+		
ajoene	+	+		+	+	+
Diallyl disulfide	+		+	+	+	+
1,5-hexa dianyl trisulfide				+		
Methyl allyl trisulfide				+		
2-vinyl 1,3- dithiene				+		
3-vinyl 1,3- dithiene				+		
S-allyl merc aptocysteine					+	
Se-methyl selenocysteine					+	
Allyl propyl disulfide			+			
Sodium 2-propenyl thiosulphinate				+		
s-methyl1- cysteine sulfoxide			+			

1996; Augusti, 1996; Block, 1998; Tahiliani & Kar, 2003; Mohamed *et al.*, 2006; Eidi *et al.*, 2006; Murugavel *et al.*, 2007). Reuter & Sendl (1995), Srivastava *et al.* (1995), Augusti (1996), Ross (1999), Warber (1999), Harris *et al.* (2001), Morbidoni *et al.* (2001), Ichikawa *et al.* (2003), Stephen (2005), Ariga & Seki (2006) and Nies *et al.* (2006) have reviewed the pharmacology and medicinal application of garlic isolated compounds for the treatment of a number of diseases (Table 1).

3.1. Antimicrobial activities

Garlic has a reputation as a cure for a variety of bacterial and fungal diseases. Louis Pasteur (1858) and Lehmann (1930) provided the first modern scientific evidences for medicinal and anti-bacterial use of garlic extract. Garlic has been observed to possess antiviral, antibacterial and antifungal activities (Cavallito & Bailey, 1944; Cavallito *et al.*, 1944; Johnson & Vaughn, 1969; Nagai, 1973; Appleton & Tansey, 1975; Tansey & Appleton, 1975; Tynecka & Gos, 1975; Sharma *et al.*, 1977; Yamada & Azuma, 1977; Amer *et al.*, 1980; Caporaso *et al.*, 1983; Elnima *et al.*, 1983; Lau *et al.*, 1983; Tutakne *et al.*, 1983; Adetumbi *et al.*, 1986; Yoshida *et al.*, 1987; Srivastava *et al.*, 1995; Lee *et al.*, 2003; Kim *et al.*, 2004; Konaklieva & Plotkin, 2006). Cavallito & Bailey (1944) had demonstrated that garlic juice diluted to one part in 125,000 inhibits the bacterial growth of *Staphylococcus*, *Streptococcus*, *Vibrio* (including *V. cholerae*) and *Bacillus* (including *Bacillus typhosus*, *B. dysenteriae* and *B. enteritidis*). Johnson & Vaughn (1969) reported that 10 % extract of dehydrated garlic bulb demonstrated antibactericidal action against *Salmonella typhimurium* and *Escherichia coli* within 2 to 6 h exposure. Crude extract of garlic has proven to be quite effective against both gram-negative and gram-positive bacteria. Bacterial cultures resistant to the commonly used antibiotic chloramphenicol appear sensitive to garlic. Jezpwa *et al.* (1966), Adetumbi & Lau (1983), Khan *et al.* (1985), Hughes & Lawson (1991), Ahsan & Islam (1996), Tsao *et al.* (2003) and Zhou (2003) observed that preparation of fresh garlic and vacuum dried powdered garlic preparations were effective against *Staphylococcus aureus*, *Streptococcus viridans*, *Streptococcus haematycticus*, *Klebsiella pneumoniae*, *Proteus vulgaris*, *Escherichia coli*, *Salmonella enteritidis*, *Bacillus subtilis*, *B. megaterium*, *B. pumilus*, *B. mycoides*, *B. thuringiensis*, *Sarcina lutea*, and *Serratia marcescens*. Al-Waili Saloom *et al.* (2007) investigated that heating, storage and ultraviolet exposure of fresh garlic juice significantly inhibited its antimicrobial activity against common human pathogens. Hughes & Lawson (1991) observed allicin, methyl allylthiosulfinates and allylmethylsulfinates, constituents in aqueous garlic clove and powder homogenate, had *in vitro* anti-bacterial activities while polar compound alliin did not. Allicin and allyl-methyl plus methyl-allyl thiosulfinate have shown inhibition of the *in vitro* growth of *Helicobacter pylori*, the bacterium responsible for serious gastric diseases as ulcer and even gastric cancer (Canizares *et al.*, 2004a,b). Sasaki & Kita (2003) reported the antibacterial activity of garlic powder against *Bacillus anthracis*. A 1% water solution of garlic powder in the test tube method killed *B. anthracis* within 3 h of treatment at room temperature. Barki & Douglas (2005) and Groppo *et al.* (2007) observed that garlic extract inhibits the growth of oral pathogens and certain proteases and thus it may have therapeutic value, particularly for periodontitis.

Anti-fungal activity of garlic extract has been demonstrated in 40 species of zoopathogenic fungi, retarding the growth in 8 of 15 genera tested (Appleton & Tansey, 1975; Shadkchan *et al.*, 2004; Ledezma & Apitz-Castro, 2006). In a comparison of the fungistatic activity of garlic extract with nystatin, griseofulvin and amphotericin B, garlic had a broad-spectrum activity against 17 strains of fungi, including the dermatophytes, yeasts, *Aspergillus* and *Penicillium*. Garlic was more effective than nystatin in retarding growth of the fungi (Srivastava, 1984). Barone & Tansey (1977) and Yamada & Azuma (1977) demonstrated that allicin in aqueous extract of garlic bulbs was the major anticandida constituent. Aqueous extract of garlic is inhibitory and lethal to numerous strains of *Cryptococcus neoformans* (Fromtling & Bulmer, 1978). In China, garlic is used frequently to treat fatal form of *Cryptococcal meningitis* (Abdullah *et al.*, 1988). Singh & Singh (1997) observed that water extract of garlic inhibits the growth of certain fungi that can cause meningitis. An aqueous extract of garlic has been demonstrated to inhibit the growth of several zoopathogenic fungi such as *Candida albicans*, commonly involved in vaginitis, *Histoplasma capsulatum*, a fungus that produces a disease similar to tuberculosis and fungi that cause athletes' foot and ringworm (Srivastava *et al.*, 1995). Yoshida *et al.* (1987), in studying the

antifungal activity of different purified fractions of garlic against *Aspergillus niger* and *Candida albicans*, have observed that ajoene has stronger activity than allicin. Adetumbi & Lau (1986) have reported that an aqueous extract of dehydrated garlic preparation inhibits the growth of the dimorphic fungus *Coccidioides immitis* and inhibits *in vitro* fungal spore germination. Adetumbi *et al.* (1986) and Lemar *et al.* (2002) reported that reduction of *Candida albicans* growth by the aqueous garlic extract is due to the inhibition of nucleic acid, protein and lipid syntheses. Stephen (2005) reported that the breakdown products of allicin, the main parent antifungal compound in garlic, have a common mode of action against antimicrobial, anticancer and anticholesterol properties. Some of the breakdown products of allicin cross cell membranes and combine with sulfur-containing molecular groups in amino acids and proteins, thus interfering with cell metabolism. Human cells are not poisoned by allicin derivatives as they contain glutathione, a sulfur-containing amino acid that combines with the allicin derivative, thus preventing cell damage (Stephen, 2005). Allicin synergized the fungicidal activity of Cu²⁺ ions against various strains of fungus (Ogita *et al.*, 2006). Allicin treatment caused Cu²⁺ solubilization complexation with a plasma membrane protein.

Nagai (1973) reported *in vivo* antiviral effect of garlic in mice against intranasally-inoculated influenza virus. The garlic extract also enhances the production of neutralizing antibody if inoculated with the influenza vaccine (Nagai, 1973). Investigation on *in vitro* activity of garlic against selected viruses, including herpes simplex virus type¹ and type², parainfluenza virus type³, vaccinia virus, vesicular stomatitis virus and human rhinovirus type², demonstrated that certain garlic constituents - (*viz.*) ajoene, allicin, allyl methyl thiosulfinate, methyl allyl thiosulfinate - are potent virucides (Weber *et al.*, 1992). Josling (2001), on the basis of his study on 146 volunteers, reported that an allicin containing garlic supplement can prevent the infection of common cold virus. Diallyl trisulfide in garlic bulb has anti-human cytomegalovirus (anti-HCMV) activity and the mechanism is associated with suppression of ie gene transcription (Zhen *et al.*, 2006).

3.2. Garlic and cardiovascular diseases

The medicinal effect of garlic and garlic extracts on cardiovascular diseases has been widely studied. Preparations of garlic and chemical constituents of garlic have been investigated for possible effects on cardiovascular diseases such as hyperlipidemia, hypertension, platelet aggregation and blood fibrinolytic activity (Kendler, 1987; Lawson *et al.*, 1992; Isensee *et al.*, 1993; Isaacsolin *et al.*, 1998; Koscielny *et al.*, 1999; Mantawy & Mahoud, 2002; Gardner *et al.*, 2003; Siegel *et al.*, 2004).

3.2.1. Hypocholesterolaemic action

Most of the studies on the effects of garlic on blood lipids in patients with cardiovascular disorders have been carried out in India. Intake of high-fat meals causes a significant increase in serum triglyceride and cholesterol levels (Groot & Scheck, 1984). Observations suggest that use of garlic (and onion as well) prevents the hypercholesterolemia induced by high-fat meal (Jain & Andleigh, 1969; Bordia & Bansal, 1973; Sainani *et al.*, 1979; Mirhadi & Singh, 1991; Elkayam *et al.*, 2003). Augusti (1977) and Bhushan *et al.* (1979) reported that eating of 10 g fresh garlic per day for 2 months significantly decreases (15%-28.5%) serum cholesterol levels among hypercholesterolemic patient. Sainani *et al.* (1979) compared vegetarians, with different eating habits with regards to the use of onions and garlic. Group I consumed onion and garlic in liberal amounts (50 g garlic and 600 g onion per week); Group II consumed small amount of onion and garlic (10 g garlic and 200 g onion per week), and Group III totally abstained from the use of garlic and onion. The level of cholesterol, triglyceride, phospholipid and β - lipoproteins were the lowest in the individuals consuming liberal amounts of garlic and onion. These results indicate that routine consumption of onion and garlic in the diet has a beneficial effect in maintaining the serum lipids at low or normal levels.

Total serum cholesterol is an important factor in the development of ischemic heart disease (IHD). Cholesterol present in the β -lipoprotein (LDL, Low density lipoprotein) and pre- β -lipoprotein (VLDL, Very low density lipoprotein) fractions finds its way into the arterial wall, whereas α -lipoprotein (HDL, High density lipoprotein) cholesterol helps to reduce the serum cholesterol level. Increase in HDL/LDL ratio is a preventive effect of the development of IHD. Jain (1977) and Bordia (1981) observed

that consumption of 0.25 mg/kg garlic oil caused a steady decrease in LDL+VLDL levels with concomitant increase in HDL levels. Ernst *et al.* (1985), Orekhov *et al.* (1995), Efendy *et al.* (1997) and Ide *et al.* (1997) have observed that the aged garlic extract, aqueous extract from garlic powder, and its constituents lower serum cholesterol level and inhibit the oxidative modification of LDL, and thus is presumed to have a protective effect against atherosclerosis.

Gebhardt (1993) reported the multiple inhibitory effects of garlic extracts at several different steps in cholesterol biosynthesis pathway in human Hep G₂ cells. According to him, defined compounds (allicin) present in water soluble extracts of garlic inhibit the biosynthesis of cholesterol in hepatocytes, thus contributing to the reduction of serum cholesterol. Garlic derived components are capable of combining with the sulphhydryl (-SH) group (Augusti & Mathew, 1974; Block, 1985). Reduced conversion of acetate into cholesterol has been observed both *in vivo* and *in vitro* (Chang & Johnson, 1980; Saxena *et al.*, 1980). As -SH groups are involved in several metabolic pathways, it may be inferred that garlic besides inhibiting lipid synthesis (Eilat *et al.*, 1996) would affect other metabolic activities as well (Srivastava *et al.*, 1995). Larner (1995) proposed the tellurium hypothesis of hypocholesterolaemic action of garlic. Accordingly, tellurium compounds found in high concentration in fresh garlic bulb may contribute to hypocholesterolaemic action by inhibiting squalene epoxidase, the penultimate enzyme in the synthetic pathway of cholesterol. Oxidation of LDL plays an important role in the initiation and progression of atherosclerosis. Yamaji *et al.* (2004) suggested that use of egg-yolk-enriched garlic powder (EGP) has antiatherogenic effect. EGP inhibits copper-induced LDL oxidation in a dose-dependent manner. Use of EGP significantly suppressed the production of peroxides in HL 60 cells and protects endothelial cells from hydrogen peroxide-induced cell injury. Sumioka *et al.* (2006) reported that *Monascus* garlic fermented extract (MGFE) attenuates hyperlipidemia suggesting that it is a potent agent for preventing arteriosclerotic diseases. Intake of enteric-coated garlic powder (equal to 400 mg garlic, 1mg allicin) twice daily has significantly reduced total cholesterol, LDL-cholesterol and triglyceride and increased HDL-cholesterol (Kojuri *et al.*, 2007).

3.2.2. Hypotensive effect

High blood pressure is one of the major risk factors of atherosclerosis (Srivastava *et al.*, 1995). The hypotensive effect of garlic was recognized by Loeper & Debray (1921). Damrau (1941) gave two Allimin tablets containing 4.75 g of garlic concentrate (equivalent to about 0.31 g of desiccated garlic and 2.375 g of desiccated parsley) to 26 hypertensive patients three times daily for three days. Blood pressure reduction was observed in 85% of the patients: the average decline in systolic and diastolic blood pressure was 12.3 mm Hg and 6.5 mm Hg, respectively. He also reported that headache was relieved in 14 of the 17 patients complaining of the symptom, and dizziness was cured in 12 of 13 patients and improvement was observed in remaining patient.

For centuries garlic has been used against hypertension in China and Japan and is recognized officially for this treatment by the Japanese food and drug administration (Bolton *et al.*, 1982). Piotrowski (1948), Papayannopoulos (1969), Srinivasan (1969) and Anonymous (1986) have also demonstrated garlic's antihypertensive effect. Pectov (1979) cited several studies on hypotensive effect of garlic from the Soviet Union and Bulgaria. Aqel *et al.* (1991) suggested that hypotensive action of garlic juice may be due to a direct relaxant effect on smooth muscles. Although components of garlic do exert pharmacological and therapeutic effects on hypertension (Srivastava *et al.*, 1995), the nature of the garlic component responsible is not known. Andrianova *et al.* (2002) observed that allisor treatment of 600 mg/day produced reduction in both systolic and diastolic blood pressure.

3.2.3. Antiplatelet aggregation activity

More recent literature suggests possible beneficial effects of garlic and its extract in preventing atherosclerotic disease. Reuter *et al.* (1974) and Lutomski (1987) demonstrated that adenosine, a compound in high concentration (0.056%) in garlic, inhibits aggregation of platelet and improves blood flow in the coronary vessels. Investigations have shown that garlic oil could inhibit platelet aggregation (Mohammad *et al.*, 1980; Mohammad & Woodward, 1986; Srivastava & Justesen, 1989) and ether extract of garlic juice taken with a fatty diet could decrease cholesterol and fibrinogen and increase

fibrinolytic activity and blood coagulation time. Garlic can lead to a decrease in thrombocyte aggregation in blood (Bordia *et al.*, 1977; Srivastava, 1984, 1986; Harenberg *et al.*, 1988; Jung *et al.*, 1989; Kiesewetter *et al.*, 1991; Lawson, 1993; Srivastava & Tyagi, 1993; MacDonald *et al.*, 2004). Srivastava & Mustafa (1993) reported that several mechanisms are involved in garlic-induced inhibition of platelet aggregation. These effects are the modification of platelet membrane properties, inhibition of calcium mobilization, and inhibition of several steps of the arachidonic acid cascade in platelets. In fact, a direct inhibitory effect of garlic extracts and its components on the enzymes of the arachidonic acid cascade have been reported (Srivastava, 1984; Srivastava & Justesen, 1989; Makheja & Bailey, 1990; Wagner *et al.*, 1991). Day *et al.* (1976) reported that several biochemical agents such as adenosine diphosphate (ADP), collagen, epinephrine, arachidonate and especially thromboxane-A elevate platelet aggregation. Bordia *et al.* (1978) observed that garlic caused a dose dependent inhibition of platelet aggregation induced by ADP, epinephrine and collagen. Both aqueous and organic extracts of garlic inhibit platelet aggregation induced by arachidonic acid, ADP, adrenaline, collagen, calcium ionophore A23187 and thrombin (Apitz-castro *et al.*, 1983; Srivastava, 1984; Srivastava & Justesen, 1989).

Makheja & Bailey (1990) reported that adenosine is one of the most important ingredients of garlic bulb. About half of the anti-platelet activity of garlic has been attributed to adenosine. They also reported that adenosine and allicin present in garlic inhibited platelet aggregation without altering arachidonic acid (AA) metabolism in platelets. Lawson *et al.* (1992) reported the antiaggregatory activity of different garlic preparations. In platelet rich plasma most of the antiaggregatory activity of garlic clove homogenates was due to adenosine; however, in whole blood the antiaggregatory activity was due to allicin and other thiosulfinates. Garlic powder tablets were equally as active as clove homogenates, whereas steam distilled oils and oil macerates only 35% and 12%, respectively. In steam distilled oils most of the activity was due to diallyl trisulfide. For the oil macerates, most of the activity was due largely to the vinylthiols. Polysulfides, such as diallyl trisulfide (Srivastava & Mustafa, 1993) and dimethyl trisulfide (Makheja & Bailey, 1990), are shown to be antiaggregatory and altered AA metabolism. Inhibition of platelet aggregation ultimately decreases atherosclerosis and thrombosis (Mirhadi & Singh, 1991). Kiesewetter *et al.* (1991) observed that intake of 800 mg of garlic powder over a period of 4 weeks caused disappearance of thrombocyte aggregation, increased microcirculation of the skin by 47.6% and decreased plasma viscosity by 3.2%; decreases in diastolic blood pressure by 9.5% and blood glucose concentration by 11.6% were also observed. Jung *et al.* (1990) investigated that single intake of 900 mg garlic powder (corresponds to 2700 mg fresh garlic) by healthy subjects caused no change in blood pressure and heart rate. Jung *et al.* (1991) reported that five hours after intake of 900mg garlic powder caused a significant increase in capillary skin perfusion by 55%. It causes minimal vasodilation and significant reduction in haematocrit and plasma viscosity. Apitz-Castro *et al.* (1983) reported that methanol extract of garlic and some pure fractions inhibited platelet aggregation. The antiplatelet property of garlic is especially due to one of its components (E) and (Z)-ajoene, which perturbs membrane by lodging itself deep between the two monolayers of plasma membranes of the intact platelets. Jain & Apitz-Castro (1987) reported that ajoene did not alter AA metabolism in intact platelets. It is of interest to note that ajoene was shown to potentiate synergistically the antiaggregatory effects of prostacyclin, forskolin, indomethacin and dipyridamole (Apitz-castro *et al.*, 1986). Chang *et al.* (2004) identified sodium 2-propenyl thiosulfinate (2PTS) from boiled garlic, which significantly inhibits ADP-induced platelet aggregation at 0.1 mM in human platelets. In contrast, the maximal aggregation percentage returned to normal level at 1mM of alk(en)yl thiosulfinate in human platelets, suggesting that 2PTS has the potential to promote immune function and prevents cardiovascular diseases. Cavagnaro *et al.* (2007) suggest that processing of garlic and the conditions used for cooking can markedly influence its effectiveness as a platelet inhibitor. They are of the opinion that crushing garlic before moderate cooking can reduce the loss of antiplatelet activity.

3.3. Anticancer potential

Garlic is one of the most ancient spice plants reputed to have an effect on cancer. As recorded around 1550 B.C. in the Ebers Papyrus, garlic was applied externally for the treatment of tumors by ancient Egyptians and internally by Hippocrates and Indian physicians (Hartwell, 1967, 1968).

Interestingly, China provides an ideal “Field Laboratory” for epidemiological studies of cancer incidence. Stomach cancer was found to rank higher for males and females in cancer mortality (Wang *et al.*, 1985; Lau *et al.*, 1990) than other cancer incidence in China (Mei *et al.* 1982). They suggested that consumption of garlic may have inhibited nitrate reduction by bacteria. Subsequently, the lower gastric nitrite (a nitrosamine precursor) concentration may reduce the risk of developing stomach cancer. You *et al.* (1989) identified that cigarette smoking, use of salty foods and moldy grains are associated with increased risk of stomach cancer. A significant reduction of stomach cancer risk was found to be associated with increasing consumption of garlic, scallions and Chinese chives (You *et al.*, 1988). Galeone *et al.* (2006) investigated the role of allium’s vegetables in the etiology of various neoplasms. They analyzed the relation between frequency of onion and garlic use and cancer at several sites of Italian and Swiss case-control studies. They reported an inverse relationship between the frequency of use of garlic and the risk of several common cancers. Abdullah *et al.* (1988) observed that natural killer-cells of raw garlic and aged group destroyed 139% and 159% more tumor cells than those from the control group. Aged garlic extract is a good chemopreventive agent for colorectal cancer. Aged garlic extract could prevent tumour formation by inhibiting angiogenesis through suppression of endothelial cell mortality, proliferation and tube formation (Matsuura *et al.*, 2006).

Aflatoxins are naturally occurring toxic metabolites produced by *Aspergillus flavus* and related fungi. Among the 16 or more toxic compounds secreted by these fungi, aflatoxin B₁ (AFB₁) is most potent in causing mutations in various microorganisms as well as mammalian and human cells. It is both toxic and hepato-carcinogenic to a variety of animals and humans in Asia and Africa (Stoloff, 1976; Wong & Hsieh, 1976; Perkin *et al.*, 1988). Tadi *et al.* (1991a,b) reported that two organosulfur compounds of garlic, ajoene and diallyl sulfide affected AFB₁ metabolism and DNA binding by inhibiting phase I enzymes and, therefore, act as potential cancer chemopreventive agents.

Garlic is one of the best natural sources of germanium. It is of interest to note that this trace metal has also been reported to prevent and cure cancer. Garlic is also an excellent source of selenium, which has potential therapeutic value in cancer treatment (Bolton *et al.*, 1982; Lawson, 1993). S-methyl selenocysteine is the major seleno-compound in garlic bulb. Epidemiological studies have indicated a negative relationship between selenium intake and the incidence of certain cancers. Blood or plasma levels of selenium are usually lower in patients with cancer than those without this disorder. In mammary tumour model, se-methyl selenocysteine was shown to be most effective seleno-compound so far in reduction of tumours (Whanger, 2004).

Although garlic has been shown to be chemopreventive against cancer, its mode of action is not fully understood. Several mechanisms have been suggested. A potential carcinogen may exert its effects by producing nuclear damage. Wargovich & Goldberg (1985) studied the effect of a sulfide and disulfide component of garlic on several chemical carcinogens. DMH (1,2-dimethylhydrazine) and NMBA (N-nitroso methyl benzylamine) are procarcinogens, which are metabolized in the liver to carcinogenic metabolites, an effect mediated by their binding to the nucleus. Pre-administration of diallyl sulfide prevented nuclear damage induced by DMH and NMBA. These observations suggest that diallyl sulfide might inhibit the conversion of procarcinogens to ultimate carcinogens, an important factor of the initiation phase of carcinogenesis. Chung *et al.* (2002) observed that garlic juice reduced endogenous formation of carcinogens N-nitrosodimethylamine (NDMA) in young human. Recently, Singh *et al.* (2006) have reported that consumption of garlic powder induced a 35-60% reduction in DNA damage induced by NDMA in rat.

Glutathione-s-transferase (GST) is an important enzyme that conjugates electrophils and assists in the detoxification of many carcinogens. Compounds that increase GST activity have been found to inhibit carcinogen induced neoplasia (Wattenberg, 1983; Sparnin *et al.*, 1988). Oral administration of methyl trisulfide and diallyl disulfide stimulated glutathione-s-transferase activity, which increased with time and showed a time lag of 18 h or more (Sparnin *et al.*, 1986; Sumiyoshi & Wargovich, 1989). These studies suggest that stimulation of glutathione-s-transferase activity in the liver and other target organs of carcinogens by garlic derived organosulphur compounds may be responsible for their protective effects in chemical carcinogenesis and on different stages of carcinogenesis.

Oxygen radicals may play a role in tumor promotion. Various anti-oxidants, trace radicals, scavengers, stimulation of super oxide dismutase, glutathione peroxidase inhibitors of cyclooxygenase and lipoxygenase enzymes were found to modulate the oxidative stress and the formation of molecular species involved in tumor promotion. Perchellet *et al.* (1986) demonstrated garlic oil stimulated glutathione peroxidase activity, and inhibited the decrease in the intracellular ratio of reduced to oxidized glutathione produced by TPA in epidermal cells. Garlic oil was also found to inhibit lipoxygenase enzymes, which is involved in TPA-stimulated arachidonic acid metabolism (Block *et al.*, 1988). Li *et al.* (1995) investigated the effect of aged garlic extract (AGE) and two of its components, s-allylcysteine (SAC) and s-allyl mercaptocysteine (SAMC), on human breast cancer cells MCF-7 and MCF-7-s. They observed an anti-proliferative response to SAC and SAMC and an alteration in glutathione level without significant concurrent changes in the glutathione metabolizing enzymes. Xiao *et al.* (2003) found that SAMC exerts anti proliferative effects by binding directly to tubulin and disrupting the microtubule assembly, thus arresting cells in mitosis and triggering JNK₁ and caspase-3 signaling pathways that lead to apoptosis.

Xu *et al.* (2004) reported a novel aspect into the understanding of the molecular mechanism of potential antitumor action of ajoene. Ajoene-treated leukemia cell line HL60 cells were arrested in G(2)/M phase and total amount of cytosolic proteasome increased. Proteasome plays a key role in the regulation of many cellular processes. Oommen *et al.* (2004) found that allicin inhibited the growth of cancer cells of murine and human origin. Allicin induced the formation of apoptotic bodies, nuclear condensation and a typical DNA ladder in cancer cells. They have observed that allicin activates the caspases-3, -8 and -9 and cleavage of poly (ADP-ribose) polymerase. Xiao *et al.* (2006) have observed that diallyl trisulfide found in garlic bulb induces apoptosis in PC-3 human prostate cancer cell by induction of protein Bax and Bak. Chu *et al.* (2006) studied the effect of garlic derivatives S-allylcysteine and S-allylmercaptocysteine on the inhibition of cancer cell invasion through restoration of E-cadherin expression. They suggest that these compounds might be potential therapeutic agents for suppressing androgen-independent prostate cancer.

It seems that garlic derived organosulfur compounds could have effects on different stages of carcinogenesis. It is now recognized that a wide range of human cancer, perhaps as many as 80-90% is attributed to environmental factors (Doll, 1977; Doll & Peto, 1987). Garlic has been shown to inhibit growth of transplantable tumors and to reduce the incidence of certain spontaneously occurring tumors. Components of garlic also inhibit the activity of diverse chemical carcinogens during both the initiation and promotion phases of carcinogenesis (Lau *et al.*, 1990; Tadi *et al.*, 1990; 1991a,b; Milner, 1996).

Kodera *et al.* (1989) isolated a new allixin (phytoalexin) synthesized by garlic. Being a phenolic compound, it is an effective inhibitor of phospholipid metabolism stimulated *in vitro* by the tumor promoter. Yamasaki *et al.* (1991) observed that allixin plays an important role in cancer prevention. Lau *et al.* (1991) and Patya *et al.* (2004) studied the effect of organosulphur compounds of garlic on the growth of animal tumors. According to Patya *et al.* (2004), allicin exhibits immune-stimulatory and antitumor properties. Immune stimulatory effect of allicin is mediated by redox-sensitive signaling such as activation of p2(ras). The antitumor effect of allicin is related to its immune-stimulatory properties. Allicin inhibits the apoptosis of macrophages in depleted nutritional state through the mitogen-activated protein kinase/extracellular signal-regulated kinase pathway (Cho *et al.*, 2006). Druesne *et al.* (2006) observed that diallyl disulfide (DADS) found in garlic bulb inhibits colon tumor cell proliferation. They suggest that 200 µM of DADS increases histone acetylation, CDKN1A mRNA and p21 (waf1) protein levels and induces G2/M cells cycle arrest in tumor cells. Agarwal *et al.* (2007) observed that pretreatment of rats with 50-100 mg garlic oil/kg body weight for a week significantly protects free radical injury and cancer in kidneys induced by ferric nitrilotriacetate (Fe-NTA). This is due to the inhibition of hepatic tumour markers viz., ornithine decarboxylase activity and DNA synthesis. It is therefore obvious that the organosulphur compounds of garlic extend their protective anti-cancer effects in three ways: (i) a direct inhibition of tumor cell metabolism; (ii) inhibition of the initiation and/or promotion phases of carcinogenesis; and (iii) modulation of the host response by augmenting macrophages and T-lymphocytes function.

3.4. Garlic as a pesticide

Effective pesticides have been developed from garlic extracts. Field trials carried out on mosquito breeding sites at Bombay (now Mumbai, India) have shown that garlic oil is very effective on several species of mosquitoes. Amonkar & Reeves (1970) studied the toxic effect of crude methanolic extract and garlic oil against 3rd stage of larvae of *Culex peus*, *C. tarsalis*, *Aedes aegypti*, *A. trisoriatus*, *A. sirensis* and 3rd and 4th stage larvae of highly insecticide resistant strains of *A. nigromaculis*. They reported that partially purified oil fraction was more toxic than the crude extract. Later, the larvicidal principles of garlic oil have been isolated and identified as diallyl disulfide and diallyl trisulfide (Amonkar & Banerji, 1971). Allicin inhibited malaria infection by inhibiting cysteine protease, which processed circumsporozoite protein (CSP) of *Plasmodium sporozoites* for invasion of host cells (Coppi *et al.*, 2006). These compounds were fatal to *Culex pipiens quinquefasciatus*. Nakagawa *et al.* (1980) reported that raw garlic juice at a dose 5 ml/kg body weight caused death of rats due to stomach injury. Gupta & Sharma (1993) have observed that root-dip treatment for tomato seedlings in 25 ppm allicin for 5 min is effective against nematode *Meloidogyne incognita*. Concentration of 200 and 100 ppm allicin as bare-root dips for 30 min killed 83% and 87% of *M. incognita* of tomato seedlings, respectively. Gareth *et al.* (2006) reported that the insecticidal activity of garlic juice in two dipterian pests *Delia radicum* (LC₅₀ - 0.4%) and *Musca domestica* (LC₅₀- 2.2%). They reported that garlic juice is effective against all the life stages of both insects.

Ajoene, the major bioactive compound derived from garlic, showed a potent leishmanicidal activity in *in vitro* against *Leishmania mexicana* and *L. donovani* (Ledezma *et al.*, 2002). The 50% inhibitory concentration (IC₅₀) for lysis was about 2 µM. They reported that leishmanicidal activity of ajoene is due to the morphological alteration of the mitochondrial membrane and nuclear envelope, as well as the formation of large autophagic vacuoles. Gamboa-Leon *et al.* (2006) suggest that garlic extract has mild protective effect against *Leishmania donovani*. This might be due to unspecific enhancement of interferon-gamma secretions. The homopteran sucking insect *Lipaphis erysimi* (mustard aphid) causes severe damage to various crops. This pest not only affects plants but it also transmits single-stranded RNA luteoviruses while feeding, which cause disease and damage in the crop. The mannose-binding garlic leaf lectin has been found to be a potent control agent of *L. erysimi* on one hand and on the other symbionin mediated luteovirus transmission (Banerjee *et al.*, 2004).

Singh & Singh (1993) reported that the water extract of garlic bulb is a potential source of molluscicide against *Lymnaea acuminata* and *Indoplanorbis exustus*. These snails are the intermediate host of liver fluke *Fasciola hepatica* and *F. gigantica*, which causes 94% fascioliasis in the buffalo's population of northern India (Singh & Agarwal, 1981; Singh & Agarwal, 1983). Singh & Singh (1995) characterized that allicin is the main molluscicidal component of garlic. The toxicity of allicin (LC₅₀ - 3.64 mg/l) at 96 h exposure is 4.1, 4.2 and 2.5 times higher with respect to synthetic molluscicides (phorate, carbaryl and formothion, respectively) (Singh & Agarwal, 1983). There was a progressive increase in the toxicity (8.67 fold) of garlic bulb extract against *Lymnaea acuminata* in the pre-harvest (2 to 5 months) period while in the post-harvest period (5 to 11 months) it was 5.5 fold (Singh & Singh, 1996a). Significant increase in toxicity of pre-harvest garlic is due to increase in level of allicin. Increase in molluscicidal activity of garlic in post harvest period is due to some changes in the chemical composition of garlic bulb (Lawson *et al.*, 1991b). Singh & Singh (1996b) demonstrated that the treatment of snail with sublethal concentrations of allicin caused a significant inhibition of acetylcholinesterase (AChE), lactic dehydrogenase (LDH) and alkaline phosphatase (ALP) activity in the nervous tissue of *Lymnaea acuminata*. Kinetic studies demonstrated that the inhibition of AChE by allicin was uncompetitive whereas inhibition of LDH and ALP was competitive (Singh & Singh, 1996b). Inhibition of this enzyme by allicin in the nervous tissue of snail *L. acuminata* may be the cause of the molluscicidal action of garlic.

Singh & Singh (1997), Singh *et al.* (1998), Singh & Singh (2000a,b), Singh & Singh (2001a,b) and Tripathi & Singh (2001) have reported that combinations of garlic bulb powder with *Cedrus deodara* oil, *Annona squamosa* seed, *Lawsonia inermis* seed, *Punica granatum* bark, *Canna indica* root powder, piperonyl butoxide and MGK-264 are very effective molluscicide. These combinations were effective in reducing the fecundity, hatchability and survival of young snails (Singh & Singh, 2004). Singh & Singh (2000b) have clearly demonstrated that these effects of garlic powder were due to inhibition of

certain enzymes and alteration in brain biogenic amine levels in the nervous tissue of *L. acuminata*. Mantawy & Mahmoud (2002) observed that glucose and glycogen levels and phenol oxidase activity in snail *Biomphalaria alexandrina* were significantly decreased after 2 and 7 day of feeding on garlic, which ultimately reduced the fecundity of snails. Spraying of garlic powder (24h LC₅₀ - 7.24%) singly and in combination with *Cedrus deodara* (24h LC₅₀ - 4.18%) were toxic, as well as effective in controlling the reproductive capacity of the giant African snail *Achatina fulica* (Rao & Singh, 2000, 2002; Rao *et al.*, 2003). Allicin, the active molluscicidal component of garlic (Singh & Singh, 1995), is effective in killing snails as well as making them sterile (Singh & Singh, 2000a).

3.5. Adverse effects

Apart from the valuable medicinal properties garlic may also act toxically when overdosed. Repeated or excessive garlic ingestion produces toxic effects (Rose *et al.*, 1990; Fehri *et al.*, 1991). The most apparent problem with using garlic in human medicine is its strong odor (Morbidoni *et al.*, 2001). A problem of clinical importance is that some people are allergic to sulfur based compounds. There were several reported allergic reactions to garlic; namely, contact dermatitis, asthma, rhinitis, conjunctivitis, urticaria, anaphylaxis and angioedema (Gaddoni *et al.*, 1994; Rauterberg, 1995; Kanerva *et al.*, 1996; Armentia & Vega, 1997; Roberge *et al.*, 1997; Morbidoni *et al.*, 2001). Consumption of garlic enhances the pharmacological effects of anticoagulants (Warfarin, fluidione) and reduces the efficacy of antiAIDS drug Saquinavir (Borrelli *et al.*, 2007).

4. Conclusion

The foregoing account emphasizes the fact that garlic is a nature's boon to mankind. A single clove of garlic has the potential of curing a man from a large number of diseases by inhibiting the population of different strains of bacteria, fungi, harmful viruses, insects and snails. Garlic's cardiovascular, antibiotic and perhaps anticancer effects are well-accepted world over because of the wealth of scientific literature supporting these effects. Use of garlic in the prevention of heart disease has been advocated by prominent researchers. Instead of several clinical tests, garlic is not yet widely recognized by medical authorities. More research should be undertaken in future to determine its efficacy as a cardiovascular preventive agent with respect to other natural product on one hand and modern drugs on the other. Anticarcinogenic components of garlic deserve urgent attention by scientist. Use of garlic as a pesticide is one of the recent concepts for eco-friendly and effective control of pest. Garlic requires investigation mainly in its mode of action in pest body and the way of delivery in the environment to ensure more effective control with less effort. Therefore, garlic deserves more attention by agriculturists and public health specialists. More investigations are still required in experimental, clinical as well as in epidemiological field to explore its full potential in the welfare of mankind.

5. Acknowledgments

The authors are grateful to Prof. Anil K. Srivastava, Former Head, of the Department of Zoology, DDU Gorakhpur University for his valuable suggestions in preparation of the manuscript.

6. References

- Abdullah TH, Kandil O, Elakdi A, Carter J. Garlic revisited: therapeutic for the major diseases of our times. *J Natn Med Ass* 1988;80:439-45.
- Adetumbi MA, Lau BHS. *Allium sativum* (garlic): a natural antibiotic. *Med Hypothesis* 1983;12:227-37.
- Adetumbi MA, Lau BHS. Inhibition of *in vitro* germination and spherulation of *Coccidioides immitis* by *Allium sativum*. *Curr Microbiol* 1986;13:73-6.
- Adetumbi MA, Javor GT, Lau BHS. Notes: *Allium sativum* (garlic) inhibits lipid synthesis by *Candida albicans*. *Antimicrobial Agents Chemother* 1986;30:499-501.
- Agarwal KC. Therapeutic actions of garlic constituents. *Med Res Rev* 1996;16:111-24.

- Agarwal MK, Iqbal M, Athar M. Garlic oil ameliorates ferric nitrilotriacetate (Fe-NTA)- induced damage and tumor promotion: implications for cancer prevention. *Food Chem Toxicol* 2007;45:1634-40.
- Ahsan M, Islam SN. Garlic: a broad spectrum antibacterial agent effective against common pathogenic bacteria. *Fitoterapia* 1996;67:374-6.
- Al-Waili Saloom KY, Akmal M, Al-waili TN, Al-waili H, Ali A, Al-Sahani K. Effects of heating, storage and ultraviolet exposure on antimicrobial activity of garlic juice. *J Med Food* 2007;10:208-12.
- Amer M, Taha M, Tassan Z. The effect of aqueous garlic extract on the growth of dermatophytes. *Int J Dermatol* 1980;19:285-7.
- Amonkar SV, Banerji A. Isolation and characteri-zation of larvicidal principle of garlic. *Science* 1971;174:1343.
- Amonkar SV, Reeves EL. Mosquito control with active principle of garlic, *Allium sativum*. *J Econ Entomol* 1970;63:1172-5.
- Andrianova IV, Fomchenkov IV, Orekhov AN. Hypotensive effect of long-acting garlic tablets allisor (a double-blind placebo-controlled trail). *Ter Arkh* 2002;74:76-88.
- Anonymous. The Wealth of India. Raw materials. New Delhi: Publication & Information Directorate, CSIR 1985;1(A):1-513.
- Anonymous. Zheziang Institute of Traditional Chinese Medicine. The effect of essential oil of garlic on hyperlipidemia and platelet aggregation. *J Tradit Chin Med* 1986;6:117-20.
- Apitz-castro R, Cabrera S, Cruz MR, Ledezma E, Jain MK. Effects of garlic extract and of three pure components isolated from it on human platelet aggregation, arachidonate metabolism, release reactivity and platelet ultrastructure. *Thromb Res* 1983;32:155-69.
- Apitz-Castro R, Escalanate J, Vargas R, Jain MK. Ajoene, the antiplatelet principle of garlic, synergistically potentiates the antiaggregatory action of prostacyclin, forskolin, indomethiacin and dypiridamole on human platelets. *Thromb Res* 1986;42:303-11.
- Appleton JA, Tansey MR. Inhibition of growth of zoopathogenic fungi by garlic extract. *Mycologia* 1975;67:882-85.
- Ariga T, Seki T. Antithrombotic and anticancer effects of garlic-derived sulfur compounds: a review. *Biofactors* 2006;26:93-103.
- Armentia A, Vega JM. Can inhalation of garlic dust cause asthma? *Allergy* 1997;51:137-38
- Aqel MB, Gharabai MN, Salhab AS. Direct relaxant effects of garlic juice on smooth and cardiac muscles. *J Ethnopharmacol* 1991;33:13-9.
- Augusti KT. Hypocholesterolemic effect of garlic (*Allium sativum* Linn.). *Indian J Exp Biol* 1977;15:489-90.
- Augusti KT. Therapeutic values of onion (*Allium cepa* L.) and garlic (*Allium sativum* L.). *Indian J Exp Biol* 1996;34:634-40.
- Augusti KT, Mathew PT. Lipid lowering effect of allicin (diallyl disulphide-oxide) on long term feeding to normal rats. *Experientia* 1974;30:468-70.
- Barki IM, Douglas CWI. Inhibitory effect of garlic extract on oral bacterial. *Arch Oral Biol* 2005;50:645-51.
- Banerjee S, Hess D, Majumder P, Roy D, Das S. The interactions of *Allium sativum* leaf agglutinin with a chaperon in group of unique receptor protein isolated from a bacterial endosymbiont of the mustard aphid. *J Biol Chem* 2004;279:23782-9.
- Barone FE, Tansey MK. Isolation, purification, identification, synthesis and kinetics of activity of the anticandidal component of *Allium sativum* and a hypothesis for its mode of action. *Mycologia* 1977;69:793-825.
- Bhushan S, Sharma SP, Singh SP, Agrawal S, Indraycin A, Seth P. Study of the hypocholesterolemic effect of onion (*Allium cepa*) on normal human beings. *Indian Med Gaz* 1979;16:249-51.
- Block E. The chemistry of garlic and onions. *Sci Am* 1985;252:114-9.
- Block E. The organosulfur chemistry of the genus *Allium* -Implication for the organic chemistry of sulfur. *Angew Chem Int Ed Engl* 1992;31:1135-78.
- Block E. Garlic as a functional food: a status report. In: Shibamoto T, Terao J, Qsawa T, editors. *Functional foods for disease prevention*. Washington: American Chemical Society, 1998. v.2, p.125-43.

- Block E, Iyer R, Grisoni S, Saha C, Belman S, Lossing FP. Lipoxygenase inhibitors from the essential oil of garlic. Markovnikov addition of the allyldithio radical to olefins. *J Am Chem Soc* 1988;110:7813-27.
- Bolton S, Null G, Troetel WM. The medical uses of garlic- fact and fiction. *Am Pharmacol* 1982;22:40-3.
- Bordia A. Effect of garlic and blood lipids in patients with coronary heart disease. *Am J Clin Nutr* 1981;34:2100-3.
- Bordia A, Bansal HC. Essential oil of garlic in prevention of atherosclerosis. *Lancet* 1973;2:1491-2.
- Bordia AK, Joshi HK, Sanadhya YK, Bhu N. Effect of essential oil of garlic on serum fibrinolytic activity in patients with coronary heart disease. *Atherosclerosis* 1977;28:155-9.
- Bordia AK, Sodhya SK, Rathore AS, Bhu N. Essential oil of garlic on blood lipids and fibrinolytic activity in patients with coronary artery disease. *J Assoc Physicians India* 1978;26:327-31.
- Borrelli F, Capasso R, Izzo AA. Garlic (*Allium sativum* L) Adverse effects and drug interactions in humans. *Mol Nutr Food Res* 2007;51:1386-97.
- Cai XJ, Uden PC, Block E, Zhang X, Quimby BD, Sullivan JJ. *Allium* chemistry: identification of natural abundance organoselenium volatiles from garlic, elephant garlic, onion and Chinese chive using headspace gas chromatography with atomic emission detection. *J Agric Food Chem* 1994;42:2081-4.
- Canizares P, Gracia I, Gomez LA, Garcia A, Martin de Argila C, Boixeda D, de Rafael L. Thermal degradation of allicin in garlic extracts and its implication on the inhibition of the *in vitro* growth of *Helicobacter pylori*. *Biotechnol Progr* 2004a;20:32-7.
- Canizares P, Gracia I, Gomez LA, Garcia A, Martin de Argila C, Boixeda D, de Rafael L. De. Allyl thiosulfinates, the bacteriostatic compounds of garlic against *Helicobacter pylori*. *Biotechnol Progr* 2004b;20:397-401.
- Caporaso N, Smith SM, Ing RHK. Antifungal activity in human urine and serum after injection of garlic. *Antimicrobial Agents Chemother* 1983;23:700-2.
- Cavagnaro PF, Camargo A, Galmarini CR, Simon PW. Effect of cooking on garlic (*Allium sativum* L.) antiplatelet activity and thiosulfinates content. *J Agric Food Chem* 2007;55:1280-8.
- Cavallito CJ, Bailey JH. Allicin: the antimicrobial principle of *Allium sativum*. I. Isolation, physical properties and antibacterial action. *J Am Chem Soc* 1944;66:1950-1.
- Cavallito CJ, Buck JS, Suter CM. Allicin: the antibacterial principle of *Allium sativum*. II. Determination of chemical structure. *J Am Chem Soc* 1944;66:1952-4.
- Chang HS, Yamato O, Sakai Y, Yamasaki M, Maede Y. Acceleration of superoxide generation in polymorphonuclear leukocytes and inhibition of platelet aggregation by alk(en)yl thiosulphates derived from onion and garlic in dogs and humans. *Prostaglandins Leukot Essent Fatty Acids* 2004;70:77-3.
- Chang MLW, Johnson MA. Effect of garlic on carbohydrate metabolism and lipid synthesis in rats. *J Nutr* 1980;112:931-8.
- Cho SJ, Rhee DK, Pyo S. Allicin a major component of garlic, inhibits apoptosis of macrophages in a depleted nutritional state. *Nutrition* 2006;22:1177-84.
- Chu Q, Ling MT, Feng H, Cheung HW, Tsao SW, Wang X, Wong YC. A novel anticancer effect of garlic derivatives: inhibition of cancer cell invasion through restoration of E-cadherin expression. *Carcinogenesis* 2006;27:2180-9.
- Chung MJ, Lee SH, Sung NJ. Inhibitory effect of whole strawberries, garlic juice or kale juice on endogenous formation of N-nitrosodimethylamine in humans. *Cancer Lett* 2002;182:1-10.
- Coppi A, Cabinian M, Mirelman D, Sinnis P. Antimalarial activity of allicin, a biologically active compound from garlic cloves. *Antimicrobial Agents Chemother* 2006;50:1731-7.
- Damrau F. The use of garlic concentrate in vascular hypertension. *Med Res* 1941;153:249-51.
- Day HJ, Holmsen H, Zucker MB. Platelet function testing. Bethesda: National Institute of Health Service, 1976. (DHEW-NIH 78-1087).
- Doll R. Strategy for detection of cancer hazards to man. *Nature* 1977;265:589-96.
- Doll R, Peto R. The cause of cancer: quantitative estimation of avoidable risk of cancer in the United States. *J Natl Cancer Inst* 1987;66:1193-8.

- Druesne PN, Pagniez A, Thomas M, Cherbuy C, Duee PH, Martel P, Chaumontet C. diallyl disulfide increases CDKN1A promoter-associated histone acetylation in human. *J Agric Food Chem* 2006;54:7503-7.
- Efendy JL, Simmons DL, Campbell GR, Campbell JH. The effect of the aged garlic extract 'Kyolic' on the development of experimental atherosclerosis. *Atherosclerosis* 1997;132:37-42.
- Eidi A, Eidi M, Esmaeili E. Antidiabetic effect of garlic (*Allium sativum* L.) in normal and streptozotocin-induced diabetic rats. *Phytomedicine* 2006;13:624-9.
- Eilat S, Avramovitch D, Rabinkov A, Mirelman D, Battler A, Elder M, Vered Z. Allicin: Hypolipidemic. *Drugs Future* 1996;21:1101-4.
- Elkayam A, Mirelman D, Peleg E, Wilchek M, Miron T, Rabinkov A, Oron HM, Rosenthal T. The effects of allicin on weight in fructose-induced hyperinsulinemic, hyperlipidemic, hypertensive rats. *Am J Hypertens* 2003;16:1053-6.
- Elnima El, Ahmed SA, Mekkawi AG, Mossa JS. The antimicrobial activity of garlic and onion extracts. *Pharmazie* 1983;38:747-8.
- Ernst E, Weihmayr TH, Matrai A. Garlic and blood lipids. *Br Med J* 1985;291:139.
- Fehri B, Aiache JM, Korbi S, Monkni M, Ben MS, Memmi A, Hizaoui B, Bouker K. Toxic effects induced by the repeated administration of *Allium sativum* L. *J Pharm Belg* 1991;46:363-74.
- Fenwick GR, Hanley AB. The genus *Allium* - Part I. *Crit Rev Food Sci Nutr* 1985;22:199-271.
- Fromtling RA, Bulmer GS. *In vitro* effect of aqueous extract of garlic (*Allium sativum*) on the growth and viability of *Cryptococcus neoformans*. *Mycologia* 1978;70:397-404.
- Gaddoni G, Selvi M, Resta F, Pasi F. Allergic contact dermatitis to garlic in a cook. *Ann Ital Dermatol Clin Sper* 1994;48:120-1.
- Galeone C, Pellucchi C, Levi F, Negri E. Onion and garlic use and human cancer. *Am J Clin Nutr* 2006;84:1027-32.
- Gamboa-Leon R, Paraguai-de-Souza E, Borja-Cabrera GP, Santos FN, Myashiro LM, Pinheiro RO, Dumonteil E, Palatanik-de-Sousa CB. Immunotherapy against visceral leishmaniasis with the nucleoside hydrolase-DNA vaccine of *Leishmania donovani*. *Vaccine* 2006;24:4863-73.
- Gardner CD, Messina M, Lawson LD, Farquhar JW. Say, garlic, ginkgo biloba: their potential role in cardiovascular diseases prevention and treatment. *Curr Atheroscler Rep* 2003;5:468-75.
- Gareth MP, Tamara SG, Andrew F. Insecticidal activity of garlic juice in two dipterian pests. *Agric For Entomol* 2006;8:1-6.
- Gebhardt R. Multiple inhibitory effects of garlic extracts on cholesterol biosynthesis in hepatocytes. *Lipids* 1993;28:613-9.
- Grosso FC, Ramacciato JC, Motta RH, Ferraresi PM, Sartoratto A. Antimicrobial activity of garlic against oral streptococci. *Int J Dent Hyg* 2007;5:109-15.
- Groot PHE, Scheck LM. Effects of fat ingestion on high density lipoprotein profiles in human sera. *J Lipid Res* 1984;4:692.
- Gupta R, Sharma NK. A study of the nematicidal activity of allicin: an active principle in garlic, *Allium sativum* L., against root-knot nematode, *Meloidogyne incognita*. (Kofoid and White 1919) Chitwood 1949. *Int J Pest Manag* 1993;39:390-2.
- Harenberg J, Giese C, Zimmermann R. Effect of dried garlic on blood coagulation, fibrinolysis, platelet aggregation and serum cholesterol levels in patient with hyperlipoproteinemia. *Atherosclerosis* 1988;74:247-9.
- Harris JC, Cottrell SL, Plummer S, Lloyd D. Antimicrobial properties of *Allium sativum* (garlic). *Appl Microbiol Biotechnol* 2001;57:282-6.
- Hartwell JL. Plants used against cancer: a survey. *Lloydia* 1967;30:379-436.
- Hartwell JL. Plants used against cancer: a survey. *Lloydia* 1968;31:71-170.
- Hughes BG, Lawson LD. Antimicrobial effects of *Allium sativum* L. (garlic), *Allium ampeloprasum* L. (elephant garlic), and *Allium cepa* L. (onion), garlic compounds and commercial garlic supplement products. *Phytother Res* 1991;5:154-8.
- Ichikawa M, Ryu K, Yoshida J, Ide N, Kodera Y, Sasaoka T, Rosen RT. Identification of six phenylpropanoids from garlic skin as major antioxidants. *J Agric Food Chem* 2003;51:7313-7.

- Ide N, Nelson AB, Lau BHH. Aged garlic extract and its constituents inhibit CU-2+- induced oxidative modification of low density lipoprotein. *Planta Med* 1997;63:263-4.
- Isaacsohn JL, Moser M, Stein EA, Dudley K, Davey JA, Liskov E. Garlic powder and plasma lipids and lipoproteins: a multi center, randomized, placebo controlled trial. *Arch Inter Med* 1998;158:1189-94.
- Isensee H, Rietz B, Jacob R. Cardioprotective actions of garlic (*Allium sativum*). *Arztl Forsch* 1993;43:94-8.
- Jain RC. Effect of garlic on serum lipids, coagulability and fibrinolytic activity. *Am J Clin Nutr* 1977;30:1380-1.
- Jain RC, Andleigh HS. Onion and blood fibrinolytic activity. *Br Med J* 1969;258:514.
- Jain MK, Apitz-castro R. Garlic: molecular basis of the putative 'Vampire repellent' action and other matters related to the heart and blood. *Trends Biochem Sci* 1987;12:252-4.
- Jezpwa J, Rafinski T, Wrocinski T. Badania nad dzialaniem antybiotycznym czosnku pospolitego (*Allium sativum* L.). *Herba Pol* 1966;12:66.
- Johnson MG, Vaughn RH. Death of *Salmonella typhimurium* and *Escherichia coli* in the presence of freshly reconstituted dehydrated garlic and onion. *Appl Microbiol* 1969;17:903-6.
- Josling P. Preventing the common cold with a garlic supplement: a double-blind, placebo-controlled survey. *Adv Ther* 2001;18:189-93.
- Jung F, Wolf S, Kiewewetter H, Mrowietz C, Pundir G, Heiden M, Miyayashita C, Wenzel E, Reim N. Wirkung von Knoblauch auf die fließfähigkeit des Blutes, Ergebnisse placebokontrollierter Pilotstudien an gesunden Probanden. *Natur und Ganzheitsmed* 1989;7:216-21.
- Jung F, Jung EM, Pundir G, Kiewewetter H. Effect of different garlic preparations on the fluidity of blood, fibrinolytic activity and peripheral microcirculation in comparison with placebo. *Planta Med* 1990;56:668.
- Jung EM, Jung F, Mrowietz C, Kiewewetter H, Pundir G, Wenzel E. Influence of garlic powder on cutaneous microcirculation. *Arzt Forschung* 1991;41:626-30.
- Kabelik J. Antimikrobielle Eigenschaften des Knoblauchs. *Pharmazie* 1970;25:266-70.
- Kanerva L, Estlander T, Jolanki R. Occupational allergic contact dermatitis from spices. *Contact Dermatitis* 1996;35:157-62.
- Kaufman PB, Pennacchini L, McKenzie M, Hoyt JE. Good and bad uses of these compounds by humans. In: Kaufman PB, Cseke LJ, Warber S, Dukes JA, Briemann HL (eds). *Natural products from plants*. New York: CRC Press, 1999. p.123-56.
- Kendler BS. Garlic (*Allium sativum*) and onion (*Allium cepa*): a review of their relationship to cardiovascular diseases. *Prev Med* 1987;16:670-85.
- Khan KI, Khan FZ, Nazar S. The antimicrobial activity at *Allium sativum* (garlic), *Allium cepa* (onion) and *Raphanus sativus* (radish). *J Pharm Pb Univ Lhr* 1985;6:59-71.
- Kiewewetter H, Jung F, Pundir G, Jung EM, Mrowietz C, Wenzel E. Effect of garlic on thrombocyte aggregation, microcirculation and risk factors. *Int J Clin Pharmacol Ther Toxicol* 1991;29:151-5.
- Kim JW, Kim YS, Kyung KH. Inhibitory activity of essential oils of garlic and onion against bacteria and yeasts. *J Food Prot* 2004;67:499-504.
- Koch HP, Lawson L. Garlic: the Science and therapeutic application of *Allium sativum* L. and related spices. 2nd ed. Baltimore: Williams & Wilkins, 1996.
- Kodera Y, Matsuura H, Yoshida S, Sumida T, Itakura Y, Fuwa T, Nishino H. Allixin, a stress compound in garlic. *Chem Pharmacol Bull* 1989;37:1656-8.
- Kojuri J, Vosoughi AR, Akrami M. Effects of *Anethum graveolens* and garlic on lipid profile in hyperlipidemic patients. *Lipids Health Dis* 2007;6:5 [doi:10.1186/1476-512 IX-6-5].
- Konaklieva MI, Plotkin BJ. Antimicrobial properties of organosulfur anti-infectives: a review of patent literature 1999-2005. *Recent Patents Anti Infect Drug Discov* 2006;1:177-80.
- Koscielny J, Klussendorf D, Latza R, Schmitt R, Radtke H, Siegel G. The antiatherosclerotic effect of *Allium sativum*. *Atherosclerosis* 1999;144:237-49.
- Larner AJ. How does garlic exert its hypocholesterolaemic action? The tellurium hypothesis. *Med Hypothesis* 1995;44:295-7.

- Lau BHS, Keeler WH, Adetumbi MA. Antifungal effect of garlic. Annu Meet Am Soc Microbiology, USA; 1983.p.387.
- Lau BHS, Tadi PP, Tosk JM. *Allium sativum* (garlic) and cancer prevention. Nutr Res 1990;10:937-48.
- Lau BHS, Yamashaki T, Gridley DS. Garlic compounds modulate macrophage and T-lymphocyte function. Mol Biother 1991;3:103-7.
- Lawson LD. Bioactive organosulfur compound of garlic and garlic products: role in reducing blood lipids. In: Kinghorn AD, Balandrin MF, editors. Human medicinal agents from plants. Washington: American Chemical Society,1993. p.306-30. (ACS Symposium Series, n. 534).
- Lawson LD. Garlic a review of its medicinal effects and indicated active compounds. In: Lawson LD, Bauer R, editors. Phytomedicines of Europe. Washington: American Chemical Society, 1998. p.176-9. (Chemistry and biological activity series, n. 691)
- Lawson LD, Block E. Comments on garlic chemistry stability of s-(2-propenyl) 2-propene-1-sulfinothioate (allicin) in blood solvents and stimulated physiological fluids. J Agric Food Chem 1997;45:542.
- Lawson LD, Hughes BG. Characterization of the formation of allicin and other thiosulfinates from garlic. Planta Med 1992;58:345-50.
- Lawson LD, Wood SG, Hughes BG. HPLC analysis of allicin and other thiosulfinates in garlic clove homogenates. Planta Med 1991a;57:263-70.
- Lawson LD, Wang ZJ, Hughes BG. Identification and HPLC quantitation of the sulfides and dialk(en)yl thiosulfinates in commercial garlic products. Planta Med 1991b;57:363-70.
- Lawson LD, Ransom DK, Hughes BG. Inhibition of whole blood platelet-aggregation by compounds in garlic clove extracts and commercial garlic products. Thromb Res 1992;65:141-56.
- Ledezma E, Apitz-Castro R. Ajoene the main active component of garlic (*Allium sativum*): a new antifungal agent. Rev Iberoam Micol 2006;23:75-80.
- Ledezma E, Jorquera A, Bendezu H, Vivas J, Perez G. Antiproliferative and leishmanicidal effect of ajoene on various Leishmania species: ultrastructural study. Parasitol Res 2002;88:748-53.
- Lee YL, Cesario T, Wang Y, Shanbrom E, Thrupp L. Antibacterial activity of vegetables and juices. Nutrition 2003;19:994-6.
- Lehmann FA. Untersuchungen uber *Allium sativum* (knoblauch). Arch Exp Pathol Pharmacol 1930;147:245-64.
- Lemar KM, Turner MP, Lloyd D. Garlic (*Allium sativum*) as an anti-candida agent: a comparison of the efficacy of fresh garlic and freeze dried extracts. J Appl Microbiol 2002;93:398-405.
- Li G, Qiao CH, Lin RI, Pinto J, Osborne MP, Tiwari RK. Anti-proliferative effects of garlic constituents in cultured human breast cancer cells. Oncol Rep 1995;2:787-91.
- Loeper M, DeBray M. Hypotensive effect of tincture of garlic. Progr Med 1921;36:391-2.
- Lutomski J. Inhalts- u. wirkstoffe einiger *Allium* Arten. Report at *Allium* Conf. Mendel-Frimmel-Gesellschaft fur Angewandte Genetic e.v.c.- Freising (FRG). 19-23 July, 1983a.
- Lutomski J. Das wichtigste uber knoblauch und knoblauch-praparate. Dtsch Apoth Ztg 1983b;123:623-4.
- Lutomski J. Components and biological properties of some *Allium* species. Poznan: Institute of the Medicinal Plants.1987. p.1-58.
- Lutomski J, Adamczewski B, Ostrowska B. Modyfikacja metody Jagera do oznaczania alliiny w czosnku i jego przetworach Lecznicyen. Farmac Polska 1968;24:575-80.
- Makheja AN, Bailey JM. Antiplatelet constituents of garlic and onion. Agents Actions 1990;29:360.
- MacDonald JA, Marchand ME, Langler RF. Improving upon the *in vitro* biological activity of antithrombotic disulfide. Blood Coagul Fibrinolysis 2004;15:447-50.
- Mantawy MM, Mahmoud AH. Effect of *Allium cepa* and *Allium sativum* feeding on glucose, glycogen, protein bands profile and phenol oxidase activity in *Biomphalaria alexandrina*. J Egypt Soc Parasitol 2002;32:271-83.
- Matsuura N, Miyamae Y, Yamane K, Nagao Y, Hamada Y, Kawaguchi N, Katsuki T, Hirata K, Suni S, Ishikawa H. Aged garlic extract inhibits angiogenesis and proliferation of colorectal carcinoma cells. J Nutr 2006;136:8425-65.

- Mei X, Wang MC, Xu HX, Pan XP, Gao CY, Han N, Fu MY. Garlic and gastric cancer- the effect of garlic on nitrite and nitrate in gastric juice. *Acta Nutr Sinica* 1982;4:53-8.
- Michahelles E. Über Neue wirkstoffe aus knoblauch (*Allium sativum*) und kuchenzwiebel (*Allium cepa* L.) [dissertation]. München: Ludwig-Maximilians Univiversität München, 1974. p.141.
- Milner JA. Garlic: its anti carcinogenic and anti-tumorigenic properties. First International Conference on East-West perspectives on functional foods. *Nutr Rev* 1996;54:582-6.
- Mirhadi SA, Singh S. Effect of garlic supplementation to cholesterol-rich diet on development of atherosclerosis in rabbits. *Indian J Exp Biol* 1991;29:162-8.
- Mohamed N, Attia T, Ali MA. Hepatoprotective activity of allicin against carbon tetrachloride induced hepatic injury in rats. *J Biol Sci* 2006;6:457-68.
- Mohammad SF, Woodward SC. Characterization of a potent inhibitor of platelet aggregation and release reaction isolated from *Allium sativum* (garlic). *Thromb Res* 1986;44:793-6.
- Mohammad SF, Brown S, Chuang HYK, Mason RG. Isolation, characterization, identification and synthesis of an inhibitor of platelet function from *Allium sativum*. *Fed Proc* 1980;39:543A.
- Morbidoni L, Arterburn JM, Young V, Mallins D, Mulrow C, Lawrence V. Garlic: its history and adverse effects. *J Herbal Pharmacother* 2001;1:63-83.
- Murugavel P, Pari L, Sitasawad SL, Kumar S, Kumar S. Cadmium induced mitochondrial injury and apoptosis in vero cells: protective effect of diallyl tetrasulfide from garlic. *Int J Biochem Cell Biol* 2007;39:161-70.
- Nagai K. Experimental studies on the preventive effect of garlic extract against infection with influenza virus. *Jpn J Infect Dis* 1973;47:321-5.
- Nakagawa S, Masamoto K, Sumiyoshi H, Kunihiro K, Fuwa T. Effect of raw and extracted-aged garlic juice on growth of young rat and their organs after preoral administration. *J Toxicol Sci* 1980;5:91.
- Nies LK, Cymbala AA, Kasten SL, Lamprecht DG, Olson KL. Complementary and alternative therapies for the management of dyslipidemia. *Ann Pharmacother* 2006;40:1984-92.
- Ogita A, Fugita K, Taniguchi M, Tanaka T. Dependence of synergistic fungicidal activity of Cu²⁺ and allicin, an allyl sulfur compound from garlic, on selective accumulation of the ion in the plasma membrane fraction via allicin-mediated phospholipid peroxidation. *Planta Med* 2006;72:875-80.
- Oommen S, Anto RJ, Srinivas G, Karungaran D. Allicin (from garlic) induced caspase-mediated apoptosis in cancer cells. *Eur J Pharmacol* 2004;485:97-103.
- Orekhov AN, Tertov VV, Sobenin IA, Pivovarova EM. Direct anti-atherosclerosis- related effects of garlic. *Ann Med* 1995;27:63-5.
- Papayannopoulos G. Garlic. *Lancet* 1969;2:962.
- Pasteur L. Memoire sur la fermentation appelle lactique. *Ann Chim Phys Ser* 1858;52:404-18.
- Patya M, Zahalka MA, Vanichkin A, Rabinkov A, Miron T, Mirelman D, Wilchek M, Lander HM, Novogrodsky A. Allicin stimulates lymphocytes and elicits an antitumor effect: a possible role of p21 ras. *Int J Immunol* 2004;16:275-81.
- Pectov U. Plants with hypotensive, antiatheromatous and coronarodilating action. *Am J Chin Med* 1979;7:197-236.
- Perchellet JP, Perchellet EM, Abney NL, Zirnstein JA, Belman S. Effects of garlic and onion oils on glutathione peroxidase activity, the ratio of reduced/oxidized glutathione and ornithine decarboxylase induction in isolated mouse epidermal cells treated with tumor promoters. *Cancer Biochem Biophys* 1986;8:299-312.
- Perkin DM, Laora E, Muir CS. "Estimates of the worldwide frequency of sixteen major cancers in 1980". *Int J Cancer* 1988;41:184-97.
- Piotrowski G. L'ail on therapeutique. *Praxix* 1948;26:488-92.
- Rao IG, Singh DK. Effect of single and binary combinations of plant derived molluscicides on reproduction and survival of the snail *Achatina fulika*. *Arch Environ Contam Toxicol* 2000;39:486-93.
- Rao IG, Singh DK. Toxic effect of single and binary treatments of synthetic and plant derived molluscicides against *Achatina fulika*. *J Appl Toxicol* 2002;22:211-5.

- Rao IG, Singh VK, Singh DK. Effect of single and binary combination of plant derived molluscicides on different enzymes activities in the nervous tissue of *Achatina fulika*. J Appl Toxicol 2003;23:19-22.
- Rauterberg A. Allergic contact dermatitis due to garlic in a cook. Aktuelle Dermatol 1995;21:317-8.
- Reuter HD, Sendl A. *Allium sativum* and *Allium ursinum*: Part 2. Pharmacology and medicinal application. Phytomedicine 1995;2:73-91.
- Reuter HD, Deninger R, Wagner H, Linker H. Investigations on the platelet aggregation inhibiting principle of garlic extract. In: Report of the 15th International Congress of the Society of Hematology;1974; Jerusalem.
- Roberge RJ, Leckey R, Spence R, Krenzelok EJ. Garlic burns of the breast. Am J Med 1997;15:548.
- Rose KD, Croissant PD, Parliament CF, Levin MB. Spontaneous spinal epidural hematoma with associated platelet dysfunction from excessive garlic ingestion: a case report. Neurosurgery 1990;26:880-2.
- Ross IA. Medicinal plants of the world chemical constituents traditional and modern medicinal uses. Totowa: Humana Press, 1999.
- Sainani GS, Desai DB, Gorhe NH, Natu SM, Pise DV, Sainani PG. Effect of dietary garlic and onion on serum lipid profile in jain community. Indian J Med Res 1979;69:776-80.
- Sasaki J, Kita J. Bacteriocidal activity of garlic powder against *Bacillus anthracis*. J Nutr Vitaminol 2003;49:297-9.
- Saxena KK, Gupta B, Kulshreshtha VK, Srivastava RK, Prasad DN. Effect of garlic treatment on isoprenaline-induced myocardial necrosis in albino rats. Indian J Physiol Pharmacol 1980;24:233-6.
- Shadkchan Y, Shemesh E, Mirelman D, Miron T, Rabinkov A, Wilchek M, Osherov N. Efficacy of allicin, the reactive molecule of garlic, in inhibiting *Aspergillus* spp. *in vitro*, and in a murine model of disseminated aspergillosis. J Antimicrob Chemother 2004;53:832-6.
- Sharma VC, Sethi MS, Kumar A, Rarotra JR. Antibacterial property of *Allium sativum* L.: *in vivo* and *in vitro* studies. Indian J Exp Biol 1977;15:466-8.
- Siegel G, Malmsten M, Pietzsch J, Schmidt A, Buddecke E, Michel F, Plock M, Schneider W. The effect of garlic on arteriosclerotic nanoplaque formation and size. J Phytomed 2004;11:24-35.
- Singh VK. Studies on molluscicidal properties of *Allium sativum* (garlic) against harmful gastropods [dissertation]. Gorakhpur: University of Gorakhpur, 1995.
- Singh O, Agarwal RA. Toxicity of certain pesticides to two economic species of snail in northern India. J Econ Entomol 1981;74:568-71.
- Singh DK, Agarwal RA. *In vivo* and *in vitro* studies on synergism with anticholinesterase pesticide in the snail *Lymnaea acuminata*. Arch Environ Contam Toxicol 1983;12:483-7.
- Singh DK, Singh A. Garlic (*Allium sativum*), a potent molluscicide. Biol Agric Hortic 1993;10:175-8.
- Singh VK, Singh DK. Characterization of allicin as a molluscicidal agent in *Allium sativum* (Garlic). Biol Agric Hortic 1995;12:119-31.
- Singh VK, Singh DK. Molluscicidal activity of pre- and post-harvest *Allium sativum* (Garlic). Biol Agric Hortic 1996a;12:311-8.
- Singh VK, Singh DK. Enzyme inhibition by allicin, the molluscicidal agents of *Allium sativum* L. (Garlic). Phytother Res 1996b;10:383-6.
- Singh K, Singh DK. Molluscicidal activity of plant derived molluscicides. J Herbs Spices Med Plants 1997;5:67-72.
- Singh K, Singh DK. Effect of different combination of MGK-264 or piperonyl butoxide with plant derived molluscicides on snail reproduction. Environ Contam Toxicol 2000a;38:182-90.
- Singh K, Singh DK. Toxicity to the snail *Lymnaea acuminata* of plant derived molluscicides in combination with synergists. Pest Manag Sci 2000b;56:889-98.
- Singh A, Singh DK. Molluscicidal activity of the custard apple (*Annona squamosa* L.) alone and in combination with other plant derived molluscicides. J Herbs Spices Med Plants 2001a;8:23-9.
- Singh A, Singh DK. Molluscicidal activity of *Lawsonia inermis* and its binary and tertiary combinations with other plant derived molluscicides. Indian J Exp Biol 2001b;39:263-8.

- Singh A, Singh DK. Effect of herbal molluscicides and their combination on the reproduction of the snail *Lymnaea acuminata*. Arch Environ Contam Toxicol 2004;46:470-7.
- Singh K, Singh A, Singh DK. The use of piperonyl butoxide and MGK-264 to improve the efficacy of some plant derived molluscicides. Pestic Sci 1998;54:145-9.
- Singh V, Belloir C, Siess MH, Le Bon AM. Inhibition of carcinogen-induced DNA damage in rat liver and colon by garlic powders with varying alliin content. Nutr Cancer 2006;55:178-84.
- Sparnins VL, Mott AW, Watenberg AW. Effects of allylmethyl trisulfide on glutathione s-transferase activity and PB-induced neoplasia in the mouse. Nutr Cancer 1986;8:211-5.
- Sparnins VL, Barany G, Wattenberg LW. Effects of organosulfur compounds from garlic and onion on benzo(a)pyrene-induced neoplasia and glutathione s-transferase activity in the mouse. Carcinogenesis 1988;9:134.
- Srinivasan V. A new antihypertensive agent. Lancet 1969;2:800.
- Srivastava KC. Effect of aqueous extract of onion, garlic and ginger on platelet aggregation and metabolism of arachidonic acid in blood vascular system: *in vitro* study. Prostaglandins Leukot Med 1984;13:227-35.
- Srivastava KC. Evidence for the mechanism by which garlic inhibits platelet aggregation. Prostaglandins Leukot Med 1986;22:313-21.
- Srivastava KC, Justesen U. Isolation and effect of some garlic components on platelet aggregation and metabolism of arachidonic acid in human blood platelets. Wien Klin Wochenschr 1989;8:293-9.
- Srivastava KC, Mustafa T. Pharmacological effects of spices: eicosanoid modulating activities and their significance in human health. Biochem Rev 1993;2:15-29.
- Srivastava KC, Tyagi OD. Effects of a garlic derived principle (ajoene) on aggregation and arachidonic acid metabolism in human blood platelets. Prostaglandins Leukot Essent Fatty Acids 1993;49:587-95.
- Srivastava KC, Bordia A, Verma SK. Garlic (*Allium sativum*) for disease prevention. South Afr J Sci 1995;91:68-77.
- Stoloff L. Occurrence of mycotoxins in food and feeds. In: Rodricks JV, editors. Mycotoxins and other fungal related food problems. Washington: American Chemical Society 1976. p.23-50.
- Stephen RD. An overview of the antifungal properties of allicin and its breakdown products: the possibility of a safe and effective antifungal prophylactic. Mycoses 2005;48:95-100.
- Stoll A, Seebeck E. Über den enzymatischen Abbau des alliiins und die Eigenschaften der Allinase. Helv Chim Acta 1949a;32:197-205.
- Stoll A, Seebeck E. Über die spezifität der allinase und die synthesen mehrerer dem Alliin verwandter verbindungen. Helv Chim Acta 1949b;32:866-77.
- Sumiyoshi H, Wargovich MA. Garlic (*Allium sativum*): a review of its relationship to cancer. Asia Pac J Pharmacol 1989;4:133-40.
- Sumioka I, Hayama M, Shimokawa Y, Shiraishi S, Tokunaga A. Lipid-lowering effect of monascus garlic fermented extract (MGFE) in hyperlipidemia subjects. Hiroshima J Med Sci 2006;55:59-64.
- Tadi PP, Robert MS, Teel W, Lau BHS. Anticandidal and anticarcinogenic potential of garlic. Int Clin Nutr Rev 1990;10:223-9.
- Tadi PP, Lau BHS, Teel RW, Herrmann CE. Binding of Aflatoxin B₁ to DNA inhibited by ajoene and diallyl sulfide. Anticancer Res 1991a;11:2037-42.
- Tadi PP, Teel RW, Lau BHS. Organosulfur compounds of garlic modulate mutagenesis, metabolism, and DNA binding of Aflatoxin B₁. Nutr Cancer 1991b;15:87-95.
- Tahiliani P, Kar A. The combined effect of *Triginella* and *Allium* extracts in the regulation of hyperthyroidism in rats. Phytomedicine 2003;10:665-8.
- Tansey MR, Appleton JA. Inhibition of fungal growth by garlic extract. Mycologia 1975;67:409-13.
- Tsao SM, Hsu CC, Yin MC. Garlic extract and two diallyl sulphides inhibits methicillin-resistant *Staphylococcus aureus* infection in BALB/cA mice. J Antimicrob Chemother 2003;52:974-80.
- Tripathi SM, Singh DK. Molluscicidal activity of *Punica granatum* and *Canna indica* combination with plant derived molluscicides against harmful snails. Malas J Appl Biol 2001;30:25-31.

- Tutakne MA, Satyanarayanan G, Bharadwaj JR, Sethi IC. Sporotrichosis treated with garlic juice: a case report. *Indian J Dermatol* 1983;28:41-5.
- Tynecka Z, Gos Z. The fungistatic activity of garlic (*Allium sativum*) *in vitro*. *Ann Univ Mariae Curie Sklodowska* 1975;30:5.
- Wagner H, Breu W, Sendl A, Steinke B. Comparative studies of extracts and pure compounds from garlic, wild garlic and onion on the arachidonic acid metabolism and platelet aggregation. In: 2nd International Symposium Pharmacy and Pharmacology and Clinical Applied *Allium sativum*; 1991 March 7-10; Berlin. p.15.
- Wang TG, You WC, Handerson BE, Blot WJ. A case control study of cancer in Shandong Province. *Natl Cancer Inst Monogr* 1985;69:9-10.
- Warber S. Modes of action at target sites. In: Kaufman PB, Cseke LJ, Warber S, Duke JA, Briemann HL, editors. *Natural products from plants*. New York: CRC Press, 1999. p.157-81.
- Wargovich MJ, Goldberg MT. Diallylsulfide: a naturally occurring thioether that inhibits carcinogen-induced nuclear damage to colon epithelial cells *in vivo*. *Mutat Res* 1985;143:127-9.
- Wattenberg LW. Inhibition of neoplasia by minor dietary constituents. *Cancer Res* 1983;43:2448-53.
- Weber ND, Andersen DO, North JA, Lawson LD, Hughes BG. *In vitro* virucidal effects of *Allium sativum* (garlic) extract and compounds. *Planta Med* 1992;58:417-23.
- Whanger PD. Selenium and its relationship to cancer: an update dagger. *Br J Nutr* 2004;91:11-28.
- Wong JJ, Hsieh DPH. Mutagenicity of Aflatoxins related to their metabolism and carcinogenic potential. *Proc Natl Acad Sci USA* 1976;73:2241-4.
- Xiao D, Pinto JT, Soh JW, Deguchi A, Gundersen GG, Palazzo AF, Shirin H, Weinstein IB. Induction of apoptosis by the garlic derived compound S-allylmercaptocysteine (SAMC) is associated with microtubule depolymerization and c-Jun NH(2)-terminal kinase 1 activation. *Cancer Res* 2003;15:6825-37.
- Xiao D, Lew KL, Kim YA, Zeng Y, Hahm ER, Dhir R, Singh SV. Diallyl trisulfide suppresses growth of PC-3 human prostate cancer xenograft *in vivo* in association with Bax and Bak induction. *Clin Cancer Res* 2006;12:6836-43.
- Xu B, Monsarrat B, Gairin JE, Girbal-Neuhauser E. Effect of ajoene, a natural antitumor small molecule, on human 20S proteasome activity *in vitro* and in human leukemic HL60 cells. *Fundam Clin Pharmacol* 2004;18:171-80.
- Yamada Y, Azuma K. Evaluation of the *in vitro* antifungal activity of allicin. *Antimicrob Agents Chemother* 1977;11:743-9.
- Yamaji K, Sarker KP, Abeyama K, Maruyama I. Antiatherogenic effects of an egg yolk-enriched garlic supplement. *Int J Food Sci Nutr* 2004;55:61-6.
- Yamasaki T, Teel RW, Lau BHS. Effect of allixin, a phytoalexin produced by garlic, on mutagenesis DNA binding and metabolism of aflatoxin B₁. *Cancer Lett* 1991;59:89-94.
- Yoshida S, Kasuga S, Hayashi N, Ushiroguchi T, Matsuura H, Nakagawa S. Antifungal activity of ajoene derived from garlic. *Appl Environ Microbiol* 1987;53:615-7.
- You WC, Blot WJ, Chang YS. Diet and high risk of stomach cancer in Shandong China. *Cancer Res* 1988;48:3518-23.
- You WC, Blot WJ, Chang YS, Ershow A, An Q, Yang JT, Hendersen BE, Fraumeni Jr JF, Wang TG. *Allium* vegetables and reduced risk of stomach cancer. *J Natl Cancer Inst* 1989;81:162-4.
- Zhen H, Fang F, Ye DY, Shu SN, Zhou YF, Dong YS, Nie XC, Li G. Experimental study on the action of allitridin against human cytomegalovirus *in vitro*: inhibitory effects on immediate early genes. *Antiviral Res* 2006;72:68-74.
- Zhou W. Acute lymphangitis treated by moxibustion with garlic in 118 cases. *J Tradit Clin Med* 2003;23:198.