



Effectiveness of the arbuscular mycorrhizal fungi use in the cherimoya (*Annona cherimola* Mill.) seedlings growth

Eficiencia del uso de hongos micorrízicos arbusculares en el crecimiento de plántulas de chirimoya (*Annona cherimola* Mill.)

William Viera^{1*}, Diego Campaña¹, Salomé Castro², Wilson Vásquez³, Pablo Viteri¹ and José Zambrano¹

¹Instituto Nacional de Investigaciones Agropecuarias (INIAP), Ecuador. ²Pontificia Universidad Católica del Ecuador, Ecuador. ³Universidad de las Américas (UDLA), Ecuador. Author for correspondence: willydnpv@hotmail.com

Rec.: 01.02.2016 Acep.: 15.07.2016

Abstract

Cherimoya (*Annona cherimola* Mill.) is native to the inter-Andean valleys of southern Ecuador and northern Peru. In Ecuador, the yield of this fruit crop is low, mainly due to agricultural management problems and poor fertilization. This research aims to assess the effect of native mycorrhizal fungi on seedling growth of cherimoya (cultivar 'Cangahua'). Sampling of soil and roots was carried out in 14 production sites of cherimoya. Soils where obtained the largest number of spores and greatest percentage of mycorrhizal colonization were those collected in Tumbabiro (plot 10), Gonzanamá, Paute and San Francisco de Atahualpa. These soils were used to propagate the inoculums in trap plants (*Sorghum vulgare*) and subsequently, they were used to inoculate the seeds and seedlings of cherimoya. There was no statistical difference to jointly analyze the results obtained in the inoculated seeds and seedlings. However, independently, the inoculum coming from the soil of Tumbabiro obtained the best results by doubling the content of total phosphorus and 47% increase in dry matter in cherimoya seedlings compared to control.

Key words: Arbuscular mycorrhizal colonization, foliar phosphorus, native mycorrhizal inoculum, trap plant.

Resumen

El chirimoyo (*Annona cherimola* Mill.), es originario de las vertientes interandinas del sur del Ecuador, norte del Perú. En el Ecuador el cultivo de chirimoya tiene un rendimiento bajo, debido principalmente a problemas de manejo agronómico y una pobre fertilización. La presente investigación tiene por objeto evaluar el efecto de las micorrizas nativas en el crecimiento de plántulas de chirimoya (cultivar 'Cangahua'). Se realizó el muestreo de los suelos y raíces en 14 sitios de producción de chirimoya. Los suelos que obtuvieron el mayor número de esporas y porcentaje de colonización de hongos micorrízicos fueron los colectados en Tumbabiro (Lote 10), Gonzanamá, Paute y San Francisco de Atahualpa. Estos suelos se utilizaron para propagar inóculo en plantas trampa (*Sorghum vulgare*) y posteriormente, se inóculos se los utilizaron para inocular semillas y plántulas de chirimoya. No existieron diferencias estadísticas al comparar los resultados de la inoculación en semillas y plántulas; sin embargo, al analizar independientemente, el inóculo proveniente del suelo de Tumbabiro obtuvo los mejores resultados al duplicar el contenido de fósforo total e incrementar en 47% la materia seca en las plántulas de chirimoya comparadas con el control.

Palabras clave: colonización micorrízica arbuscular, fósforo foliar, inóculo micorrízico nativo, planta trampa.

Introduction

The cherimoya (*Annona cherimola* Mill.), has its center of origin in the southern part of Ecuador, northern Peru and Bolivia (Van Zonneveld *et al.*, 2012). This fruit is widely appreciated for its taste, aroma and nutrient content. In Ecuador, this crop is mainly located in the provinces of Loja, Azuay, Pichincha, Imbabura and Tungurahua, in the Inter-Andean Region of Ecuador, at altitudes ranging from 1500 to 2600 m. a. s. l. According to the National Institute of Statistics and Census of Ecuador, the country has a total of 16201 trees of this fruit in the whole country, with a production of 447 t, which is low compared to the great demand for consumption of this fruit.

Most species of fruit trees require a period of growth in the nursery prior to transplantation in commercial orchards. In this stage, the use of mycorrhizal-arbuscular fungi show great potential because they play an important role in the growth and nutrition of higher plants, such as fruit species (Vázquez *et al.*, 2011).

These fungi are obligate symbionts and need to be associated with the root of their hosts to obtain carbon from the photosynthesis. In exchange they return to the plant the nutrients and water that are extracted from the soil. In addition, they represent a prolongation of the roots of the plant beyond the nutrient depletion zone (Saravesi *et al.*, 2014). According to the above, mycorrhizal fungi facilitate the development of cherimoya seedlings due to increased absorption of nutrients, mainly phosphorus, an element responsible for the development of new cells and thus plant growth.

The benefits of inoculation with fungi that form arbuscular mycorrhizal are expressed in enhanced seedling survival; increased plant growth in less time, thus reducing time spent in the nursery, savings in fertilizer costs; and increases in production and product quality (Verbruggen *et al.*, 2013). The modification of the root system by symbiotic association with mycorrhizal fungi helps to improve the absorption and transport of water and nutrients from the soil to the root through the increase in the volume of soil explored, which is reflected in the improvement in plant growth (Verbruggen *et al.*, 2013).

In agriculture, mycorrhizal fungi deserve special interest for their effect on plant growth through increasing the efficiency of the root system in the absorption of nutrients, especially phosphorus, and promoting absorption of trace elements (Smith *et al.*, 2015). Furthermore, the use of these fungi reduce the application of pesticides because mycorrhizal create physical barriers or activate defense mechanisms against

root pathogenic organisms associate with various metabolic routes (Goicoechea *et al.*, 2013), thus becoming an environmentally friendly alternative that helps to preserve the environment and promotes an approach to new technologies orientated towards sustainable management that can be included in the process of integrated fruit production (Verbruggen *et al.*, 2013). According to this background. The aim of this research was to evaluate the effect of mycorrhizal fungi on the growth promotion and development of seedlings of cherimoya cultivar 'Cangahua' in the nursery.

Materials and methods

Study area. The research was conducted in the laboratory (23 °C and 41% relative humidity) and in the greenhouse (29 °C and 37% relative humidity) of the Tumbaco Experimental Farm belonging to the National Institute of Agricultural Research, located at Tumbaco in the Province of Pichincha, County of Quito, at an altitude of 2348 m. a. s. l., latitude 00° 13' 00" South and longitude 78° 24' 00" West.

Mycorrhizal content analysis of the soil from cherimoya orchards

Collection of biological material. Roots and soil samples were taken in cherimoya producing areas in the Highlands of Ecuador (Table 1).

Soil sampling. To determine the number of spores in soil, sampling was carried out around the crown (dripping zone) of cherimoya plants, selected at random, at a depth of 20 cm. The sample was composed of 10 sub-samples of 100 g each one. They were transported using a cooler to the laboratory and they were dried at environmental temperature (22 °C), under shade, for 15 days.

Root sampling. This sampling was carried out to estimate the root colonization. The treetop extension was used as a reference for taking root samples. A shovel was used to dig to a depth of 40 cm around the tree at the dripping zone in order to obtain secondary and tertiary, healthy, young and slightly lignified roots. Samples were transported using a cooler to the laboratory.

Estimated variables. Mycorrhizal colonization rate in roots: The clarification and root staining method, mycorrhizal spore population in soil: spore quantification was carried out following the described method of Phillips & Hayman, (1970).

Data analysis. An analysis of variance was carried out. In addition, the Tukey statistical test at 5% to determine significance ranges. The statistical analysis was established in R 3.3.1. software™. Soils (Tumbabiro-Lote 10, San

Table 1. Soil sampled in cherimoya producing areas in Ecuador

Soil	Province	County	Parish	Site	Altitude (m.a.s.l.)	Latitude	Longitude
s1	Imbabura	Urcuquí	Tumbabiro	Lote 5	2120	00° 28' 3" N	78° 11' 0"W
s2	Azuay	Cuenca	Paute	Tutucan	2289	2° 46' 39" S	79° 45' 32" W
s3	Tungurahua	Patate	Los andes	Lote 5	2038	01°14'18"S	78°30'04"W
s4	Loja	Gonzanamá	Gonzanamá	El Tablón	2042	4° 13' 49" S	79° 29' 52" W
s5	Tungurahua	Patate	Los andes	Lote 10	2038	01°14'18"S	78°30'04"W
s6	Pichincha	Quito	Atahualpa	San José de Minas	1980	0° 09' 58" N	78° 23' 10"W
s7	Pichincha	Quito	Puéllaro	Munango	2060	0° 37' 10"N	78° 24' 44"W
s8	Pichincha	Quito	Perucho	Cementerio	2761	0° 66' 46" N	78° 22' 65" W
s9	Azuay	Cuenca	Gualaceo	Nallig		03°19'47"S	79°03'28"W
s10	Imbabura	Urcuquí	Tumbabiro	Lote 10	2120	0° 28' 3" N	78° 11' 0"W
s11	Pichincha	Quito	Atahualpa	San José de Atahualpa	2042	0° 82' 04" N	72° 22' 79"W
s12	Pichincha	Quito	Atahualpa	San Francisco de Atahualpa	2100	0° 79' 65" N	78° 22' 13" W
s13	Loja	Loja	Vilcabamba	Mollepamba	2046	04° 08' 44" S	79° 24'30"W
s14	Imbabura	Cotacachi	Panvachupa	San Pedro	2449	0° 18' 55" N	78° 15' 39" W

Francisco de Atahualpa, Paute & Gonzanamá), which showed the best results in number of spores and root colonization (Table 2), were selected to continue the assay in the next phase.

Table 2. Root colonization by mycorrhizal fungi and spores associated to soil from cherimoya producing areas in Ecuador

Soil	Origin	Numbers of spores.100g ⁻¹ soil	Colonization percentage (%)
s1	Tumbabiro Lote 5	2000cde	8,74 a
s2	Paute	2730 b	5,08 abc
s3	Patate Lote 5	1510efg	1,47 bc
s4	Gonzanamá	2770 ab	7,51 abc
s5	Patate Lote 10	1700def	0,74 c
s6	San José de Minas	1710def	1,84 abc
s7	Munango	1035 g	4,36 abc
s8	Perucho	1675def	0,52 c
s9	Gualaceo	1645ef	0,86 bc
s10	Tumbabiro Lote 10	3220 a	7,82 ab
s11	San José de Atahualpa	1170fg	2,40 abc
s12	San Francisco de Atahualpa	2410bc	5,21 abc
s13	Vilcabamba	2225bcd	3,48 abc
s14	Cotacachi	1585efg	4,11 abc
	Average	1956.07 ± 637.62	3.92 ± 3.37

Propagation in trap plants (*Sorghum vulgare*)

Soil sampling for physical-chemical analysis.

Following the sample procedure described previously, 1 k of soil sample from the four selected soils for evaluation in the trap plants was collected, labeled and sent to the laboratory for examination.

Treatments. Soils that achieved the best averages in spore count and high root colonization, constituted the inoculum that was propagated in trap plants of sorghum variety 'INIAP 201', the treatments being as follow: s1= Tumbabiro, s2= San Francisco Atahualpa, s3= Paute, s4= Gonzanamá and s5= control (sterile sand).

Experimental design and statistical analysis.

A randomized complete block design with three replications was used. In addition, Tukey test at 5% was done to determine differences between treatments. The experimental unit consisted of a pot of 2 k of capacity with 500 g of sterile sand placed in the bottom, then 1 kg of soil (treatment) and a layer of sterile sand (100 g). Ten seeds of sorghum were sown in each pot. The statistical analysis was carried out in R software version 3.3.1. TM.

Estimated variables. Total phosphorus concentration in plant tissue: After 3 months, total phosphorus concentration in plant tissue was determined by chemical composition analysis. Percentage of dry matter produced: After 90 days, the fresh weight of root, stem

and leaves was measured. Samples were placed in an oven for 24 hours at 110 °C to determine the dry weight and establish their percentage ratio. Population of mycorrhizal spores in the soil: Quantification of spores was carried out 90 days after the seeding of the trap plants, using the sieving and centrifugation method (Phillips & Hayman, 1970). Mycorrhizal colonization rate in roots: This was evaluated 90 days after the seeding of trap plants, using the method of clarification and staining roots, following the technique of Phillips & Hayman (1970), together with the method of visual density.

Specific management of the experiment in trap plants. Sorghum seeds were placed in rubbing alcohol for 3 minutes, washed with a solution of chlorine at 1.5% for 2 minutes and rinsed five times with sterile distilled water. In each pot was placed 500 g of sterile sand (water vapor at 80 °C for 1 hour), 1 kg of soil sampled and finally a layer (100 g) of sterile sand, and ten seeds were planted. At 90 days after planting, three plants at random were taken and root colonization was assessed. Using 20 g of the sieved soil, spores determination by sieving and centrifugation method was performed. Three other plants were used for determining the percentage of dry matter. The four remaining plants were maintained to continue generating the inoculum that was used in the nursery stage. Ten days before using the inoculum, the trap plants were cut by the stem base and watering was stopped. The inoculum was prepared by mixing the roots of the trap plants, which were cut into pieces of about 1 cm, with the potting soil.

Evaluation in cherimoya seeds and seedlings

Once propagated the inoculum of each of the four selected soils was assessed in nursery conditions. This phase was divided into two trials where the inoculum was evaluated in cherimoya (cultivar 'Cangahua') seeds and seedlings.

Treatments. These were constituted by four inoculums, a commercial control and an absolute control: i1= Tumbabiro, i2= San Francisco de Atahualpa, i3= Paute, i4= Gonzanamá, i5= commercial inoculum; i6= control (black soil, pomina and rice husk, all sterilized).

Evaluated variables. Rate of mycorrhizal colonization in roots: At the fourth month after the inoculation of the cherimoya seeds and seedlings (3 months old), the rate of colonization was determined by the technique of Phillips and Hayman (1970). Dry matter percentage: At 120 days after inoculation of seeds and seedlings, the fresh weight of root, stem and leaves was measured in plants, and then samples were placed in an oven for 24 hours at 110 °C to measure

the dry weight and calculate the percentage of dry matter. Total phosphorus: After 90 days, the total phosphorus was estimated by foliar analysis using the molybdenum-vanadate colorimetric method, while the molybdenum blue method was applied for phosphorus determination in the soil (AOAC, 1995).

Experimental design and statistical analysis. A randomized complete block with five replications was performed. In addition, Tukey test at 5% was estimated to determine differences among the means of treatments. The experimental unit consisted of a seedling cherimoya planted in a bag containing 2 kg of substrate with 100 g of inoculum. The statistical analysis was established in R software version 3.3.1.™.

Specific management of cherimoya seeds and seedlings. Seed disinfection: Seeds of cultivar 'Cangahua' were placed in rubbing alcohol for 3 minutes, then were soaked with a solution of chlorine at 1.5% for 2 minutes and rinsed five times with sterile distilled water. Seed pre-germination: Seeds were immersed in a solution of gibberellic acid at a concentration of 1250 ppm for 24 hours. Sowing: In plastic bags (2 k of capacity), 1.5 k of sterile substrate (black soil, pomina and rice husk in proportion 2: 1: 1) and 100 g of the inoculum was placed in each treatment. The substrate was sterilized using water vapor at 80 °C for 1 hour. One seed of cherimoya was sown and covered by 100 g more of the sterile substrate. The seedlings were previously germinated and after 90 days of growing were transplanted into the bags. Fertilization: This was done using ammonium nitrate at a dose of 2 g.l⁻¹ of water.

Results and discussion

Mycorrhizal content analysis of the soil

According to the results, there is a moderately acceptable correlation (0.64) between the number of spores and the percentage of colonization. Soils that showed the highest values in these two variables corresponded to samples collected in Tumbabiro-Lote 10 (Imbabura), Gonzanamá (Loja), Paute (Azuay) and San Francisco de Atahualpa (Pichincha) (Table 2), being in these soils the genus *Glomus* as the most frequent. The best results were recorded in the samples from Tumbabiro with a sandy loam texture (Table 3), which allows good drainage, as mentioned also by López *et al.* (2015), who stated that there is a greater presence of these microorganisms in well-drained soils. Moreover, mycorrhizal fungi act in the soil structure by providing more aggregation and stability, and preventing erosion processes (Lozano *et al.*, 2015).

Table 3. Physical-chemical characteristics of selected soils used as inoculum in the greenhouse and nursery phase

Soil	Origin	Texture	Humidity (%)	Density	pH	Matter Organic (%)
s10	Tumbabiro Lote 10	franco - arenoso	26.2	1.46	7	4.3
s12	San Francisco de Atahualpa	Franco	32.9	1.42	7.4	4.5
s2	Paute	franco - arcilloso	29	1.32	7.2	4
s4	Gonzanamá	franco - arcilloso	36	1.23	6.6	8.6

Propagation in trap plants

Dry matter percentage. When analyzing this variable (Table 4), it can be shown mycorrhizal treatments showed the best results, consistent with those reported by Bressan *et al.* (2001), who found mycorrhizal fungi contribute to increase dry matter. The inoculum produced with the soil from Gonzanamá obtained the highest value, with an average percentage of 65.58% of biomass, whereas the control without inoculation obtained 33.54%. Díaz *et al.* (2014), mention that in *S. vulgare*, it has been observed that mycorrhizal plants have increased growth, photosynthetic pigments and stomatal conductance compared with plants without mycorrhizal inoculation. Apparently the carbohydrate flow is regulated by the host plant and depends on the species of fungus.

Table 4. Variables estimated in trap plants inoculated with the different soils

Soil	Origin	Variables			
		Dry Matter (%)	Total Phosphorus (%)	Numbers of spores /100g of soil	Colonization (%)
s1	Tumbabiro-Lote 10	52.20 b	0.28 a	3247 a	21.55 a
s2	San Francisco de Atahualpa	52.12 b	0.29 a	2487 b	17.68 a b
s3	Paute	41.22 c	0.23 b	1521 c	13.60 c
s4	Gonzanamá	65.58 a	0.17 c	2771 b	12.18 c
s5	Testigo	33.54 c	0.12 d	40 d	3.12 d
	Average	48.93 ± 12.19	0.22 ± 0.07	2013.20 ± 1270.51	13.63 ± 6.92

Total phosphorus. The highest percentage of phosphorus was observed in plants inoculated with soil from Tumbabiro and San Francisco de Atahualpa with an average rate of 0.28% and 0.29%, respectively; while the control (without inoculation) obtained a much lower value (0.12%) (Table 4). Smith *et al.* (2015), reported that mycorrhizal fungi contribute to greater absorption of N, P, K, Zn and Cu by the plant, a result also

confirmed by the research done by Díaz *et al.*, (2014).

Mycorrhizal spores population in soil. The largest population of spores was observed in the treatment with the soil from Tumbabiro with an average of 3 247 spores.100 g⁻¹ of soil, while the lowest percentage was found in the control with 40 spores.100g⁻¹ of soil (Table 4). Pérez (2011), mentions that compacted soils reduce soil fertility and distribution of plant roots and, therefore, the hyphae of arbuscular-mycorrhizal fungus in the rhizosphere decrease. In addition, the formation and function of mycorrhizal fungus may be affected, finding the soil environment can favor the development of fungi at one time and reduce it at another either by direct effects on fungal communities or indirectly through the effect of the soil environment on the host plant.

Rate of mycorrhizal colonization in roots. The highest percentage of root colonization was observed in the treatment with the soil from Tumbabiro, with an average of 21.55%; while the lowest percentage (3.12%) was shown by the control (Table 4). According to these results, it is deduced that the kind of spores of the Tumbabiro soil were effective in the trap plants evaluated because the infection rate exceeded 20%, being a positive result as mentioned by Herrera (1993), which it is important because the inoculum is from the original soil without increment of the population prior to inoculation. Colonization in control plants occurred due to the capacity of survival of mycorrhizal spores in the soil, although the substrate was disinfected.

Evaluation in cherimoya seeds and seedlings

Dry matter percentage. As in the previous case, no statistical difference (p-value = 0.32) was found between seed and seedling inoculation. Seedlings with inoculum obtained from soils of Tumbabiro and Gonzanamá showed the highest percentages of dry matter compared to the other treatments (Table 5). This benefit could be explained by the colonization and action of mycorrhizal fungi. Posada & Franco (2006), reported the interaction of rhizosphere microorganisms with mycorrhizal fungi, which allows the symbiotic association to produce a greater effect on colonization, and therefore in the production of dry matter since absorption and nutrient availability is greater for the plant. Furthermore Saravesi *et al.* (2014), report the synthesized carbohydrates in leaves, have allowed the development of mycorrhizal fungi. These carbohydrates are more developed in inoculated seedlings than in seeds because seedlings start with a certain amount of leaves and are capable of carrying out photosynthesis, benefiting the fungus with carbohydrates.

Table 5. Variables estimated at 120 days after inoculation of cherimoya seeds and seedlings


Inoculum origin	Variables					
	Total Phosphorus (%)		Dry Matter (%)		Colonization (%)	
	Seed	Seedling	Seed	Seedling	Seed	Seedling
Tumbabiro-Lote 10	0.15 a	0.18 ab	23.56 b	24.83 a	7.22 a	6.04 a
San Francisco de Atahualpa	0.13 a	0.17 b	21.14 bc	18.78 bc	6.16 ab	3.83 b
Paute	0.08 bc	0.18 ab	19.10 c	19.20 bc	2.93 c	5.42 a
Gonzanamá	0.14 a	0.19 a	29.54 a	21.92 ab	6.29 ab	5.51 a
Comercial control	0.10 b	0.07 c	19.76 bc	21.04 abc	5.34 b	3.64 bc
Control (without inoculation)	0.07 c	0.07 c	18.76 c	16.98 c	2.31 c	2.72 c
Average	0.11 ± 0.03	0.14 ± 0.06	21.98 ± 4.10±	20.46 ± 2.76	5.04 ± 1.98	4.53 ± 1.31

Total phosphorus. There was no statistical difference (p -value = 0.14) when the treatment means were analyzed jointly in the percentages obtained in seed and seedling inoculation. Independently, both seeds and seedlings showed statistical differences among treatments. The largest proportion of absorbed total phosphorus was observed in the seedlings planted with the inoculum from Tumbabiro & Gonzanamá (Table 5), implying that this occurred due to the action of the mycorrhizal fungus that perform their function through external hyphae that increase the number of absorption sites (Noda, 2009). In addition, Meinhardt & Gehring (2012), found there is a greater absorption of phosphorus because mycorrhizal fungi change the pH of the soil by organic agents and solubilizers such as phosphatase, making available phosphorus which is retained in the soil, as occurred in our study (Table 6).

Table 6. Percentage of phosphorus retained in the soil before and after mycorrhizal inoculation in cherimoya seeds and seedlings

Origin	Phenological stage	Retained phosphorus - Before (%)	Retained phosphorus - After (%)
Tumbabiro-Lote 10	Seed	94.5	73.1
	Seedling	94.5	78.8
San Francisco de Atahualpa	Seed	94.5	75.8
	Seedling	94.5	75.4
Paute	Seed	94.5	71.5
	Seedling	94.5	79.1
Gonzanamá	Seed	94.5	78.8
	Seedling	94.5	73.7
Comercial control	Seed	94.5	79.8
	Seedling	94.5	75.4
Control (without inoculation)	Seed	94.5	80.2
	Seedling	94.5	79.1

Studies have shown the phosphorus concentration in plant tissue is related to the structure that colonizes the root, namely, when the root is colonized by arbuscules, the amount of phosphorus in tissue decreases because fungi also absorb phosphorus for growth; whereas phosphorus becomes available to the plant when the root is colonized by vesicles and mycelium as occurred in this research (Figure 1), and often vesicles act as a reservoir (Vanhove & Van Damme, 2013)

**Figure 1.** Vesicles and hyphae observed in secondary roots of cherimoya seedlings. The arrow indicates the vesicle.

Rate of mycorrhizal colonization in roots.

There was no statistically significant difference (p -value = 0.5) in colonization rate when comparing jointly the treatment means of the inoculation in seeds and seedlings. This is consistent with those results reported by Carreón *et al.* (2014), who stated the inoculation of these fungi can be performed in seedbeds or in transplantation. Inoculation in seedbeds could present certain problems because seeds can germinate slowly or, when they germinate, may produce certain allelopathic compounds that inhibit the process of intra-radical colonization by mycorrhizal fungi. Moreover, in most tree fruit species, seedlings depend on the reserve content of their cotyledon leaves, thus inoculation might be needless (Carreón *et al.*, 2014).

In seedling, the highest percentage of colonization (Table 5), was obtained by the treatment in which the inoculum from Tumbabiro (7.22%) was used. This percentage is apparently low because of the short time of assessment (4 months) between inoculation and the colonization analysis, taking into account that the initial growth of the root and vegetative development are slow. Control showed a percentage of colonization of 2.72%,

it can be explained by the persistence of spores in the soil due to populations of mycorrhizal fungi are composed of spores of different ages and in distinct states of dormancy or quiescence (Montañez *et al.*, 2010). Studies show in perennial species more time is required to obtain higher and steady colonization percentages because fruit crops such as cherimoya have an extended life cycle which can reach more than 30 years of life (Posada & Franco, 2006).

Conclusion

Soils sampled in Tumbabiro and Gonzanamá (main cherimoya producing areas) showed the highest populations of mycorrhizal spores and colonization rates in cherimoya roots. There was a 100% increase in the absorption of phosphorus compared to the control (without inoculation) in the trap plants where these soils were used as inoculum. Cherimoya seedlings inoculated with inoculum made with soil from Tumbabiro showed the highest colonization of roots, dry matter and accumulated phosphorus in the plant tissue. These results suggest that the use of mycorrhizal fungus is an alternative to improve the nutrient absorption, mainly phosphorus, and increase dry matter, which is reflected in the seedling development.

Acknowledgment

The authors thanks to the National Institute of Agricultural Research (INIAP) for funding this research. In addition, thanks to Lyn Jackson for her assistance in the English edition of this manuscript.

References

AOAC-Association of Official Analytical Chemist. (1995). *Official Methods of analysis* (16th. Ed.). Washington, USA: A.O.A.C. p745.

Bressan, W., Siqueira, J., Vasconcellos, C., & Purcino, A. (2001). Mycorrhizal fungi and phosphorus on growth, yield and nutrition of intercropped grain sorghum and soybean. *Pesqui Agropecu Bras*, 36 (1), 315-323. <http://dx.doi.org/10.1590/S0100-204X2001000200015>

Carreón, Y., Aguirre, S., Gavito, M., Mendoza, D., Juárez-Chávez, R., Martínez-Trujillo, M., & Trejo Aguilar, D. (2014). Inoculación micorrizica arbuscular en portainjertos de plantas de aguacate cv 'Hass' en viveros de Michoacán, México. *Rev Mex Cienc Agric*, 5(5), 847-857.

Díaz, A., Salinas, J., Espinoza, F., Garza, M., Grageda, O. (2014). Características de planta, suelo y productividad entre sorgo fertilizado e inoculado con micorriza arbuscular. *Rev Mex Cienc Agric*, 5 (3), 379-390.

Goicoechea, N., Garmendia, I., Sánchez, M., Aguirreolea, J. (2013). Arbuscular mycorrhizal fungi (AMF) as bioprotector agents against wilt induced by *Verticillium* spp. in pepper: a review. *Span J Agric Res*, 8(S1), 25-42. <http://dx.doi.org/10.5424/sjar/201008S1-5300>

Herrera, R. (1993). General methodology to analyze rootlets raw humus and vesiculo arbusculares mycorrhizal (VAM) components. La Habana, Cuba: Ed. Ancoras. p 85.

López, B., Alarcón, A., Quintero, R., Lara, A. (2015). Selección de cepas de hongos micorrizicos arbusculares en dos sistemas de producción de chile. *Rev Mex Cienc Agric*, 6(6), 1203-1214.

Lozano, J., Armbrecht, I., Montoya, J. (2015). Hongos formadores de micorrizas arbusculares y su efecto sobre la estructura de los suelos en fincas con manejos agroecológicos e intensivos. *Acta Agron*, 64(4), 289-296. <https://doi.org/10.15446/acag.v64n4.46045>

Meinhardt, K., & Gehring, C. (2012). Disrupting mycorrhizal mutualisms: a potential mechanism by which exotic tamarisk outcompetes native cottonwoods. *Ecol Appl*, 22(2), 532-549. <http://dx.doi.org/10.2307/41416780>

Montañez, I., Vargas, C., Cabezas, M., & Cuervo, J. (2010). Colonización micorrizica en plantas de aguacate (*Persea americana* L.). *Rev U.D.C.A Actual & Divulg Cient*, 13, 51-60.

Noda, Y. (2009). Las Micorrizas: Una alternativa de fertilización ecológica en los pastos. *Pastos y Forrajes*. 32 (1), 1-10.

Pérez, A., Rojas, J., & Montes, V. (2011). Hongos formadores de micorrizas arbusculares: una alternativa biológica para la sostenibilidad de los agroecosistemas de praderas en el caribe colombiano. *Rev Colomb Cienc Anim*, 3 (2), 366-385.

Phillips, J., & Hayman, D. (1970). Improved procedures for clearing roots and staining parasitic and vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. *Trans Br Mycol Soc*, 55(1), IN16-IN18. [http://dx.doi.org/10.1016/S0007-1536\(70\)80110-3](http://dx.doi.org/10.1016/S0007-1536(70)80110-3)

Posada, R., & Franco, L. (2006). Tiempo de establecimiento de pasturas y su relación con la micorriza arbuscular en paisajes de Loma y Vega. *Acta Biol Colomb*, 11 (1), 55-64.

Saravesi, K., Ruotsalainen, A., & Cahill, J. (2014). Contrasting impacts of defoliation on root colonization by arbuscular mycorrhizal and dark septate endophytic fungi of *Medicago sativa*. *Mycorrhiza*, 24(4), 239-245. <http://dx.doi.org/10.1007/s00572-013-0536-5>

Smith, S., Anderson, I., Smith, F. (2015). Mycorrhizal associations and phosphorus acquisition: from cells to ecosystems. *Ann Plant Rev* 48, 409-440. <http://dx.doi.org/10.1002/9781118958841.ch14>

Van-Zonneveld, M., Scheldeman, X., Escribano, P., Viruel, M., Van Damme, P., García, W., Tapia, C., Romero, J., Siguéñas, M., & Hormaza, J. (2012). Mapping genetic diversity of cherimoya (*Annona cherimola* Mill.): application of spatial analysis for conservation and use of plant genetic resources. *PLoS One*, 7(1), 1-14. <http://dx.doi.org/10.1371/journal.pone.0029845>

Vanhove, W., & Van Damme, P. (2013). Value Chains of cherimoya (*Annona cherimola* Mill.) in a centre of diversity and its on-farm conservation implications. *Trop Conserv Sci*, 6 (2), 158-180.

Vázquez, M., Arévalo, L., Jaen, D., Escamilla, J., Mora, A., Hernández, E., & Téliz, D. (2011). Effect of *Glomus mosseae* and *Entrophospora colombiana* on plant growth, production, and fruit quality of "Maradol" papaya (*Carica papaya* L.). *Sci Hortic*, 128(3), 255-260. <http://dx.doi.org/10.1016/j.scienta.2011.01.031>

Verbruggen, E., Heijden, M., Rillig, M., & Kiers, E. (2013). Mycorrhizal fungal establishment in agricultural soils: factors determining inoculation success. *New Phytol*, 197(4), 1104-1109. <http://dx.doi.org/10.1111/j.1469-8137.2012.04348.x>