

**FEEDING HABITS OF THE AMERICAN CROCODILE,
CROCODYLUS ACUTUS (CUVIER, 1807)
(REPTILIA: CROCODYLIDAE)
IN THE SOUTHERN COAST OF QUINTANA ROO, MEXICO**

Alejandro VILLEGAS¹ y Juan Jacobo SCHMITTER-SOTO

El Colegio de la Frontera Sur, Av. Centenario km 5.5, 77900, Chetumal, Quintana Roo,
MÉXICO

¹Present address: Universidad Autónoma Metropolitana, Unidad Xochimilco, Calzada del
Hueso 1100, Col. Villa Quietud, Delegación Coyoacán, C.P. 04960, MÉXICO, D.F.
avillegas@universo.com

RESUMEN

Entre enero y septiembre de 2006 se capturaron 21 cocodrilos (*Crocodylus acutus*) y se recapturaron seis en cuatro sistemas lagunares de la costa sur de Quintana Roo para analizar los contenidos estomacales. Las presas encontradas en el contenido estomacal se dividieron en cinco grupos y se realizaron análisis basados en la frecuencia y en el volumen. Las presas encontradas en orden de importancia fueron peces, crustáceos, aves, mamíferos e insectos. La dieta de los cocodrilos cambió considerablemente de acuerdo con el tamaño corporal de los individuos. Los cocodrilos adultos agregaron a su dieta presas de mayor talla como aves y mamíferos. Peces y crustáceos estuvieron representados en la dieta de individuos de todas las clases de tamaño, demostrando la importancia que estas presas tienen en la dieta de *C. acutus*.

Palabras clave: *Crocodylus acutus*, Quintana Roo, contenido estomacal, hábitos alimentarios.

ABSTRACT

Between January and September 2006, 21 crocodiles (*Crocodylus acutus*) were captured and six were recaptured in four lagoons located in the southern coast of Quintana Roo. Stomach contents were obtained in order to identify prey items, which were analyzed by the frequency and volume method. In decreasing order of importance, prey items were fish, crustaceans, birds, mammals and insects. Diet of crocodiles changed considerably according to their body size. Adult crocodiles added larger preys (birds and mammals) to their diet; fish and crustaceans were present in the diet at all sizes, thus showing the importance of these prey in the diet of American crocodiles.

Key words: *Crocodylus acutus*, Quintana Roo, stomach contents, feeding habits.

INTRODUCTION

The American crocodile (*Crocodylus acutus*) is distributed along the Atlantic and Pacific coasts, from Mexico to South America, the Caribbean Islands (Cuba, Jamaica, and Hispaniola), and southeastern Florida in the United States (Thorbjarnarson 1989). Diet represents an important ecological feature of every organism since it affects the body condition, behavior, growth, and reproduction

(Rice 2004). Crocodiles ingest a wide variety of prey as a result of large ontogenetic change in their body size (Barr 1997). In order to understand the ecology and behavior of crocodiles in their natural habitat, it is important to have information on the feeding habits and how these might vary with season, body size, sex and habitat. This paper documents the feeding habits of American crocodiles in several small coastal lagoons in the southern Caribbean coast of Mexico. The study focuses on diet changes according to location, season, ontogeny, and sex.

MATERIALS AND METHODS

Crocodiles were captured from four lagoons in southern Quintana Roo (Mexican Caribbean, Fig. 1), during the first five days of each month (January-September 2006), either directly by hand or by means of a pole with nose wire, depending on size. Once on board (a 3.6 m aluminum boat with a 15 HP outboard engine), sex and total length (TL) of crocodiles were determined. Crocodiles were classified into four size classes: hatchlings (TL<30 cm), juveniles (TL=30-90 cm), subadults (TL=90-180 cm), and adults (TL>180 cm) (Platt & Thorbjarnarson 2000). Animals were marked by clipping tail scutes in a coded pattern (Platt & Thorbjarnarson 1997). The hose and Heimlich maneuver technique described by Fitzgerald (1989) was used to extract all the stomach contents, PVC tubes were used to keep the jaws open and a hose of appropriate size (five to 15 mm diameter) coated with vegetable oil was inserted into the stomach; then water was introduced until stomach was completely full. A light massage in the abdominal region (Heimlich maneuver) resulted in the expulsion of the stomach content. Stomach contents were preserved in 10% formalin to stop digestion and later stored in 70% ethanol.

Fishes were identified according to Castro-Aguirre *et al.* (1999) and Schmitter-Soto (1998), while identification of other groups followed Howell and Webb (1995) for birds, Villalobos (1998) and Williams (1984) for crustaceans, and Domínguez (1990) and Morón *et al.* (1997) for insects. In addition, the help of experts was sought and the biological collection of ECOSUR was consulted for reference.

Prey found in the stomach content were categorized as fresh (recently ingested) or old (not recently ingested), because the chitinous exoskeleton of insects and crustaceans can remain in the stomach of crocodiles for several months (Delany & Abercrombie 1986, Thorbjarnarson 1993, Barr 1997).

Dietary niche breadth was measured by sex and size class using the equation of Levins (1968, cited by Krebs 1999). Kruskal-Wallis and χ^2 tests were used to look for significant differences ($P < 0.05$) in the relative volume of prey items by size class, sex, and location. The c^2 was used to test the null hypothesis of uniform distribution of data.

RESULTS

A total of 21 crocodiles (12 females, nine males) were captured comprising subadults (76%, $n=16$), adults (14%, $n=3$) and juveniles (9%, $n=2$). Six individuals were recaptured once, and the minimum time to recapture was one month in only two individuals. Night surveys were conducted starting 20-30 min after sunset to ensure sufficient darkness to detect eye shine (Messel *et al.* 1981), and finishing at dawn. With the exception of one individual captured in a small sandy beach, most crocodiles were captured next or under the roots of mangroves (*Rhizophora mangle*) along the shores of lagoons. One female (154 cm TL) had an empty stomach. The survey in Laguna Xcalak was complete, as well as in Laguna Cementerio. In Río Huach a survey of the whole lagoon system was not feasible because in some sites access was too difficult. In Bacalar Chico the only survey conducted was in some channels.




Figure 1. Localization of lagoons where individuals of *Crocodylus acutus* were captured in southern Quintana Roo.

Table 1. Percent volume of food items in stomach contents of *Crocodylus acutus* in southern Quintana Roo.

LT (cm)	Sex	Date 2006	Fish	Crustaceans	Birds	Insects	Mammals
80	Female	Apr 5	100	-	-	-	-
96.5	Female	Jul 11	87	13	-	-	-
100	Male	Jan 18	84	16	-	-	-
100	Male	Aug 17	-	98	-	2	-
106	Female	Aug 16	88	9	-	3	-
107	Female	Jul 12	99	1	-	-	-
124	Male	Jan 18	100	-	-	-	-
134	Female	Feb 27	100	-	-	-	-
140	Female	Jun 13	-	100	-	-	-
142	Male	Apr 4	74	-	19	7	-
142	Female	Apr 5	92	-	-	-	8
154	Female	Apr 20	-	-	-	-	-
168	Female	Jun 13	-	100	-	-	-
179	Male	Sep 13	52	48	-	-	-
187	Male	Jul 10	100	-	-	-	-
200	Male	Mar 1	100	-	-	-	-
203	Male	Sep 13	60	49	-	0.2	35
210	Male	May 16	72	28	-	-	-
210	Male	Jun 14	-	-	100	-	-

Stomach contents. Five prey groups were found in the stomach contents. Females prefer fish and crustaceans, in that order. Males have a more diversified diet, composed of fish, mammals, birds, crustaceans and insects (Tables 1, 2). Kruskal-Wallis test showed no significant differences ($H = 5.78$, $P = 0.016$), excepting the relationship between birds consumed by each sex; only males preyed upon birds.

Four fish taxa were identified based on remains of scales (*Gerres cinereus*, *Cyprinodon artifrons*, *Gambusia yucatanana*, and *Eucinostomus* sp.). Crustacean remains were mainly appendages, and species identification was very difficult, because it is usually necessary to have at least part of the carapace. Three crustacean species (*Panoplax depressa*, *Palaemonetes intermedius* and *Callinectes* sp.) were identified. As for mammals, only hair masses with connective tissue were found and

Table 2. Percent volume of food items in stomach contents of captured (C) and recaptured (R) *Crocodylus acutus* in southern Quintana Roo.

LT (cm)	Sex	Date 2006		Fish		Crustaceans		Birds		Insects		Mammals	
		C	R	C	R	C	R	C	R	C	R	C	R
		87.5	Female	Jan 18	May 16	67	80	33	-	-	-	-	20
116	Female	Jul 12	Sep 12	99	50	1	-	-	-	-	50	-	-
125	Female	Aug 16	Sep 12	-	-	99	20	-	-	1	0.1	-	79
185	Male	Apr 4	May 15	-	-	-	-	100	100	-	-	-	-

species identification was not possible. Remains of a beetle of the family Hydrophilidae and five individuals of the genus *Heterosternus* were found. Also, remains of one individual of the order Hemiptera and one of the family Stratiomyidae were found. There were bird remains belonging to three species (*Phalacrocorax brasilianus*, *P. auritus* and *Anas discors*) in four different individuals of *C. acutus*.

Eleven food items were obtained in the dry season and 14 in the rainy season. During the rainy season food consumption was higher (60.15% total volume) than during the dry season (39.85%), but the main prey item in both seasons was fish.

Juvenile and subadult crocodiles consumed mostly fish, followed by crustaceans and insects although the latter also included mammals in their diet. Adult crocodiles consumed mainly mammals and birds as well as some crustaceans, fish and insects.

Dietary niche breadth was significantly greater ($\chi^2 = 11.55$; $df = 2$; $P < 0.001$) in subadult crocodiles (BA = 0.085) than in juveniles (BA = 0.027) or adults (BA = 0.032).

Crocodiles from Laguna Cementerio had greater food volume in their stomachs ($H = 11.17$, $P = 0.010$) compared to individuals from the remaining sites. The χ^2 goodness-of-fit test indicated that food relative volume was not uniform among prey types ($\chi^2 = 13.66$; $df = 5$; $P < 0.001$).

DISCUSSION

Stomach contents composition of *C. acutus* was very similar to that previously reported by Gómez (2004) in Sian Ka'an Biosphere Reserve, north of our study area. However, variation in stomach contents is more apparent in its biomass, attributable to differences in fresh and old prey proportions in our samples. A high proportion of fresh prey in the samples indicates that the crocodiles are feeding frequently (Rice 2004). Analysis of stomach contents in this study suggested that fresh fish were usually consumed by crocodiles, as fish remains together with scales were found in the majority of stomachs. Gómez (2004) reported only fish scales in *C. acutus* stomach contents, and although these represented an old food item, it was however the most important item in the crocodiles' diet.

Alvarez del Toro (1974) and Thorbjarnarson (1989) stated that this crocodile shows piscivorous habits. In contrast, Casas-Andreu & Barrios (2003) found that the main prey of *C. acutus* in the Pacific coast of Mexico were mammals, birds and fish. Other authors have also reported fish as the main component of the diet for crocodylians (Delany & Abercrombie 1986, Thorbjarnarson 1993, Tucker *et al.* 1996). Differences on diet might depend on the type of habitat (Delany & Abercrombie 1986, Magnusson *et al.* 1987), but we suspect that findings of Casas-Andreu & Barrios (2003) are biased by the method used (feces analysis instead of stomach contents).

Fish diversity is relatively high in the wetland systems located within the study area, despite their limited size (Avilés-Torres *et al.* 2001). Crocodiles seem to be eating the most likely available small fish species. Large fish (lutjanids, haemulids or albulids) were absent from the stomach contents, even though they are very abundant in these lagoons. These wetlands are a functional unit that includes different biotopes (coral reefs, mangroves, seagrasses and coastal lagoons), which are utilized by a huge diversity of species in different life stages (larvae, juveniles, adults) for breeding, feeding, or reproduction (Claro 1994).

In this study, fishes were the main prey for juvenile and subadult crocodiles; whereas mammals and birds were for adults. These differences might be a consequence of skull (jaw) development, thus providing an increase in the possibility of diet diversification, since it gives the capacity of consuming larger prey (Tucker *et al.* 1996). Concerning the ingestion of birds, males consume them more than females, which may be related to the fact that males were larger (TL) than females. Similarly, ontogenetic changes on diet reflect the ability of larger individuals to catch larger preys (Magnuson *et al.* 1987, Fitzgerald 1989).

Gómez (2004) did not find insects in the stomach contents of *C. acutus*. This prey item was found to be minimally ingested by adults in our study area, although probably insects are accidental or secondarily ingested. Delany & Abercrombie (1986), Barr (1997) and Rice (2004) pointed out that ingestion of invertebrates increases with the size of crocodiles. Occurrence of large prey on diet of crocodiles enormously contributes to biomass, since they prefer them to maximize nutritional efficiency when different preys have the same availability (Wolfe *et al.* 1987). Crustaceans were an abundant prey for subadult crocodiles in our study area, especially mangrove crabs and crabs of genus *Brachyura*, similar to the findings of Gómez (2004). Blue crab (*Callinectes sapidus*) is very common in the lagoons, so this prey item seems to be consumed opportunistically by *C. acutus*. Subadults had a broader niche as they consumed five prey types.

Less than a half (44%) of the analyzed stomach contents contained vegetation, mainly marine grass and small pieces of wood. Several authors have also found plant remains in the stomachs of crocodylians considering these to be accidentally ingested

(Valentine *et al.* 1972, Delany & Abercrombie 1986, Thorbjarnarson 1989, 1993, Mohd & Stuebing 1996, Casas-Andreu & Barrios 2003).

Acknowledgments. We thank CONACYT and ECOSUR for their support to this project. To John Thorbjarnarson and David González for comments to the manuscript, and Rogelio Cedeño and Roberto Herrera for providing logistic support throughout the study. Field work was ably assisted by René Calderón, Erika Pérez, Roberto Rojo, Rodolfo Cisneros, Brenda Amenityro, Ariane Dor, Arturo Olascoaga, Axa Molina, Ferenc Worum, Martín Marrufo, Ana Lilia Díaz and Mitzi Olvera. Secretaría de Medio Ambiente y Recursos Naturales (SEMARNAT) granted permit. We thank two reviewers who provided helpful comments on the manuscript.

LITERATURE CITED

- Allstead, J., & C. Vaughan.** 1994. Food habits of *Caiman crocodilus* in Caño Negro, Costa Rica. *Vida Silvestre Neotropical* 3:24-29.
- Alvarez del Toro, M.** 1974. *Los Crocodylia de México (Estudio Comparativo)*. Instituto Mexicano de Recursos Naturales Renovables, México.
- Avilés-Torres, S., J. J. Schmitter-Soto & R. Barrientos M.** 2001. Patrones espaciales de la riqueza de peces en lagunas costeras del sur de Quintana Roo, México. *Revista de Hidrobiología*. 11:141-148.
- Barr, B. R.** 1997. Food habits of the American Alligator, *Alligator mississippiensis*, in southern Everglades. Ph.D. Dissertation, Universidad of Florida, Gainesville. Unpublished.
- Casas-Andreu, G. & Q. G. Barrios.** 2003. Hábitos alimenticios de *Crocodylus acutus* (Reptilia: Crocodylidae) determinados por el análisis de sus excretas en la costa de Jalisco, México. *Anales Instituto de Biología, UNAM, Serie Zoología* 74: 35-42.
- Castro-Aguirre, J. L., H. Espinosa & J. J. Schmitter-Soto.** 1989. *Ictiofauna estuarino-lagunar y vicaria de México*. Noriega-Limusa e Instituto Politécnico Nacional, México.
- Claro, R. (ed.)**. 1994. *Ecología de los peces marinos de Cuba*. Centro de Investigaciones de Quintana Roo, Chetumal, México.
- Delany, F. M. & C. L. Abercrombie.** 1986. American alligator food habits in northcentral Florida. *Journal of Wildlife Management* 50:348-353.
- Domínguez, R. R.** 1990. *Taxonomía: Strepsiptera a Hymenoptera, claves y diagnosis*. Universidad Autónoma Chapingo, México.
- Fitzgerald, L. A.,** 1989. An evaluation of stomach flushing techniques for crocodylians. *Journal of Herpetology* 23:170-172.
- Gómez, Y. H.** 2004. Hábitos alimentarios en el cocodrilo de río (*Crocodylus acutus*) y en el cocodrilo de pantano (*Crocodylus moreletii*) en la Reserva de la Biosfera de Sian Ka'an, Q. Roo, México. Informe Final de Servicio Social, Universidad Autónoma Metropolitana, Unidad Xochimilco, México. Unpublished.
- Howell, S. N. & S. Webb.** 1995. *A Guide to the Birds of Mexico and Northern Central America*. Oxford University Press, New York.
- Krebs, C. J.** 1999. *Ecological Methodology*. Addison-Wesley, Menlo Park.
- Kushlan, J. A. & F. J. Mazzotti.** 1989. Population biology of the American Crocodile. *Journal of Herpetology* 23:7-21.
- Magnusson, W. E., E. Vieira Da Silva & A. L. Lima.** 1987. Diets of Amazonian crocodylians. *Journal of Herpetology* 21:85-95.
- Messel, H., G. C. Vorlicek, A. G. Wells & W.J. Green.** 1981. *Surveys of Tidal River Systems in the Northern Territory of Australia and their Crocodile Populations*. Pergamon Press, Sydney, Australia.

- Mohd, S. A. S. & R. B. Stuebing.** 1996. Diet, growth and movements of juvenile crocodiles *Crocodylus porosus* Schneider in the Klias River, Sabah, Malaysia. *Journal of Tropical Ecology* 12:651-662.
- Morón, M. A., B. C. Ratcliffe & C. Deloya.** 1997. *Atlas de los escarabajos de México. Coleoptera: Lammellicornia. Vol I. Familia Melolonthidae. Subfamilias Rutelinae, Dynastinae, Cetoniinae, Trichinae, Vaginae y Melolonthidae.* Sociedad Mexicana de Entomología A. C. y la Comisión Nacional para el Conocimiento y uso de la Biodiversidad (CONABIO), México.
- Platt, S. G. & J. B. Thorbjarnarson.** 1997. Status and life history of the American crocodile in Belize. Belize Coastal Zone Management Project BZE/92/G31. Report to the United Nations (Unpubl). Development Program, Global Environment Facility, Belmopan, Belize.
- _____. 2000. Status and conservation of the American Crocodile, *Crocodylus acutus*, in Belize. *Biological Conservation*. 96:13-20.
- Rice, A. N.** 2004. Diet and condition of American alligators (*Alligator mississippiensis*) in three central Florida lakes. Master's Thesis, Universidad of Florida, Gainesville. Unpublished.
- Ross, J. P. (ed.).** 1998. *Crocodiles. Status Survey and Conservation Action Plan.* 2a ed. IUCN/SSC Crocodiles Specialist Group, Switzerland.
- Schmitter-Soto, J. J.** 1998. *Catálogo de los peces continentales de Quintana Roo.* ECOSUR, San Cristóbal de las Casas, México.
- Thorbjarnarson, J. B.** 1989. Ecology of the American crocodile, *Crocodylus acutus*. In: Crocodiles. Their ecology, management and conservation. Pp. 228-259. In: *IUCN Publication New Series, Switzerland.*
- _____. 1993. Diet of the spectacled caiman (*Caiman crocodilus*) in the central Venezuelan llanos. *Herpetologica* 49:108-107.
- Tucker, A. D., C. J. Limpus, H. I. McCallum & K. R. McDonald.** 1996. Ontogenetic dietary partitioning by *Crocodylus johnstoni* during the dry season. *Copeia* 1996:978-988.
- Valentine, J. M., J. R. Walther, K. M. McCartney & L. M. Ivy.** 1972. Alligator diets on the Sabine National Wildlife Refuge, Louisiana. *Journal of Wildlife Management*. 3:809-815.
- Villalobos, J. J. H.** 1998. *Taxonomía de los crustáceos con énfasis en el orden Decapoda.* Consejo Nacional de Ciencia y Tecnología (CONACYT) y El Colegio de la Frontera Sur (ECOSUR), México.
- Williams, A. B.** 1984. *Shrimps, lobsters, and crabs of the Atlantic coast of the Eastern United States, Maine to Florida.* Smithsonian Institution, Washington, D. C.
- Wolfe, J. L., D. K. Bradshaw & R. H. Chabrek.** 1987. Alligator feeding habits: new data, a review. *Northeast Gulf Science* 9:1-8.

Recibido: 23 de noviembre de 2007

Aceptado: 20 de mayo de 2008